




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07D 471/04, 487/04, A61K 31/435, 31/505 // (C07D 487/04, 239:00, 209:00) (C07D 471/04, 221:00, 209:00)</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/51600 (43) International Publication Date: 14 October 1999 (14.10.99)</p>
<p>(21) International Application Number: PCT/US99/07254 (22) International Filing Date: 1 April 1999 (01.04.99) (30) Priority Data: 60/080,451 2 April 1998 (02.04.98) US (71) Applicant (for all designated States except US): NEUROGEN CORPORATION [US/US]; 35 Northeast Industrial Road, Branford, CT 06405 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): HORVATH, Raymond, F. [US/US]; 239B Twin Lakes Road, North Branford, CT 06471 (US). DARROW, James, W. [US/US]; 4 Dinatale Drive, Wallingford, CT 06492 (US). MAYNARD, George, D. [US/US]; 27 Glenwood Road, Clinton, CT 06413 (US). (74) Agent: SARUSSI, Steven, J.; McDonnell Boehnen Hulbert & Berghoff, Suite 3200, 300 South Wacker Drive, Chicago, IL 60606 (US).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published With international search report.</p>
<p>(54) Title: AMINOALKYL SUBSTITUTED 9H-PYRIDINO[2,3-b]INDOLE AND 9H-PYRIMIDINO[4,5-b]INDOLE DERIVATIVES</p>		
<p>(57) Abstract</p> <p>Disclosed are compounds of formula (I), wherein Ar, R¹, W and X are substituents as defined herein, which compounds are (1) antagonists at CRF₁ receptors and are, therefore, useful in the diagnosis and treatment of stress related disorders such as post traumatic stress disorder (PTSD) as well as depression, headache and anxiety; and (2) are neuropeptide Y₁ receptor antagonists, and are therefore useful in the treatment of a variety of clinical conditions which are characterized by the presence of an excess of neuropeptide Y.</p> <div style="text-align: center;">  <p>(I)</p> </div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Aminoalkyl Substituted 9H-Pyridino[2,3-b]indole and 9H-Pyrimidino[4,5-b]indole Derivatives

BACKGROUND OF THE INVENTION

5 Field of the Invention

The present invention relates to aminoalkyl substituted 9H-pyridino[2,3-b]indole and 9H-pyrimidino[4,5-b]indole derivatives which selectively bind to corticotropin-releasing factor (CRF₁) receptors and to mammalian neuropeptide Y (NPY₁) receptors. It further relates to pharmaceutical compositions containing such compounds and the use of such compounds in treating physiological disorders induced or facilitated by CRF or those associated with an excess of neuropeptide Y.

15 Description of the Related Art

Corticotropin releasing factor (herein referred to as CRF), a 41 amino acid peptide, is the primary physiological regulator of proopiomelanocortin (POMC) derived peptide secretion from the anterior pituitary gland [J. Rivier et al., 20 *Proc. Nat. Acad. Sci. (USA)* 80:4851 (1983); W. Vale et al., *Science* 213:1394 (1981)]. In addition to its endocrine role at the pituitary gland, immunohistochemical localization of CRF has demonstrated that the hormone has a broad extrahypothalamic distribution in the central nervous system and produces a wide 25 spectrum of autonomic, electrophysiological and behavioral effects consistent with a neurotransmitter or neuromodulator role in brain [W. Vale et al., *Rec. Prog. Horm. Res.* 39:245 (1983); G.F. Koob, *Persp. Behav. Med.* 2:39 (1985); E.B. De Souza et al., *J. Neurosci.* 5:3189 (1985)]. There is also 30 evidence that CRF plays a significant role in integrating the response of the immune system to physiological, psychological, and immunological stressors [J.E. Blalock, *Physiological Reviews* 69:1 (1989); J.E. Morley, *Life Sci.* 41:527 (1987)].

Clinical data provide evidence that CRF has a role in psychiatric disorders and neurological diseases including depression, anxiety-related disorders and feeding disorders. A role for CRF has also been postulated in the etiology and pathophysiology of Alzheimer's disease, Parkinson's disease, Huntington's disease, progressive supranuclear palsy and amyotrophic lateral sclerosis as they relate to the dysfunction of CRF neurons in the central nervous system [for review see E.B. De Souza, *Hosp. Practice* 23:59 (1988)].

10 In affective disorder, or major depression, the concentration of CRF is significantly increased in the cerebral spinal fluid (CSF) of drug-free individuals [C.B. Nemeroff et al., *Science* 226:1342 (1984); C.M. Banki et al., *Am. J. Psychiatry* 144:873 (1987); R.D. France et al., *Biol. Psychiatry* 28:86 (1988); M. Arato et al., *Biol Psychiatry* 25:355 (1989). Furthermore, the density of CRF receptors is significantly decreased in the frontal cortex of suicide victims, consistent with a hypersecretion of CRF [C.B. Nemeroff et al., *Arch. Gen. Psychiatry* 45:577 (1988)]. In addition, 20 there is a blunted adrenocorticotropin (ACTH) response to CRF (i.v. administered) observed in depressed patients [P.W. Gold et al., *Am J. Psychiatry* 141:619 (1984); F. Holsboer et al., *Psychoneuroendocrinology* 9:147(1984); P.W. Gold et al., *New Eng. J. Med.* 314:1129 (1986)]. Preclinical studies in rats and 25 non-human primates provide additional support for the hypothesis that hypersecretion of CRF may be involved in the symptoms seen in human depression [R.M. Sapolsky, *Arch. Gen. Psychiatry* 46:1047 (1989)]. There is preliminary evidence that tricyclic antidepressants can alter CRF levels and thus 30 modulate the numbers of CRF receptors in brain [Grigoriadis et al., *Neuropsychopharmacology* 2:53 (1989)].

There has also been a role postulated for CRF in the etiology of anxiety-related disorders. CRF produces anxiogenic effects in animals and interactions between benzodiazepine /

non-benzodiazepine anxiolytics and CRF have been demonstrated in a variety of behavioral anxiety models [D.R. Britton et al., *Life Sci.* 31:363 (1982); C.W. Berridge and A.J. Dunn *Regul. Peptides* 16:83 (1986)]. Preliminary studies using the putative
5 CRF receptor antagonist α -helical ovine CRF (9-41) in a variety of behavioral paradigms demonstrate that the antagonist produces "anxiolytic-like" effects that are qualitatively similar to the benzodiazepines [C.W. Berridge and A.J. Dunn *Horm. Behav.* 21:393 (1987), *Brain Research Reviews* 15:71
10 (1990)]. Neurochemical, endocrine and receptor binding studies have all demonstrated interactions between CRF and benzodiazepine anxiolytics providing further evidence for the involvement of CRF in these disorders. Chlordiazepoxide attenuates the "anxiogenic" effects of CRF in both the conflict
15 test [K.T. Britton et al., *Psychopharmacology* 86:170 (1985); K.T. Britton et al., *Psychopharmacology* 94:306 (1988)] and in the acoustic startle test [N.R. Swerdlow et al., *Psychopharmacology* 88:147 (1986)] in rats. The benzodiazepine receptor antagonist (Ro 15-1788), which was without behavioral
20 activity alone in the operant conflict test, reversed the effects of CRF in a dose-dependent manner while the benzodiazepine inverse agonist (FG 7142) enhanced the actions of CRF [K.T. Britton et al., *Psychopharmacology* 94:306 (1988)].

It has been further postulated that CRF has a role in
25 immunological, cardiovascular or heart-related diseases such as hypertension, tachycardia and congestive heart failure, stroke and osteoporosis. CRF has also been implicated in premature birth, psychosocial dwarfism, stress-induced fever, ulcer, diarrhea, post-operative ileus and colonic hypersensitivity
30 associated with psychopathological disturbance and stress.

The mechanisms and sites of action through which the standard anxiolytics and antidepressants produce their therapeutic effects remain to be elucidated. It has been hypothesized however, that they are involved in the


suppression of the CRF hypersecretion that is observed in these disorders. Of particular interest is that preliminary studies examining the effects of a CRF receptor antagonist (α -helical CRF ₉₋₄₁) in a variety of behavioral paradigms have demonstrated that the CRF antagonist produces "anxiolytic-like" effects qualitatively similar to the benzodiazepines [for review see G.F. Koob and K.T. Britton, In: *Corticotropin-Releasing Factor: Basic and Clinical Studies of a Neuropeptide*, E. B. De Souza and C .B Nemeroff eds., CRC Press p221 (1990).]

Neuropeptide Y, a peptide first isolated in 1982, is widely distributed in the central and peripheral neurons and is responsible for a multitude of biological effects in the brain and the periphery. Various animal studies have shown that activation of neuropeptide NPY₁ receptors is related to vasoconstriction, Wahlestedt et al., *Regul. Peptides*, 13: 307-318 (1986), McCauley and Westfall, *J. Pharmacol. Exp. Ther.* 261: 863-868 (1992), and Grundemar et al., *Br. J. Pharmacol.* 105: 45-50 (1992); and to stimulation of consummatory behavior, Flood and Morley, *Peptides*, 10: 963-966 (1989), Leibowitz and Alexander, *Peptides*, 12: 1251-1260 (1991), and Stanley et al., *Peptides*, 13: 581-587 (1992). Grundemar and Hakanson, *TIPS*, 15: 153-159 (1994), state that, in animals, neuropeptide Y is a powerful stimuli of food intake, and an inducer of vasoconstriction leading to hypertension. They further point out that low levels of neuropeptide Y is associated with loss of appetite. These reports clearly indicate that compounds that inhibit the activity of this protein will reduce hypertension and appetite in animals.

SUMMARY OF THE INVENTION

This invention provides novel compounds of Formula I which interact with CRF₁ receptors and NPY₁ receptors. It further relates to the use of such compounds, pharmaceutical compositions comprising these compounds, and methods useful for the treatment of psychiatric and affective disorders and neurological diseases, including major depression, headaches, anxiety-related disorders, post-traumatic stress disorder, supranuclear palsy, as well as treatment of immunological, cardiovascular or heart-related diseases, hypertension, feeding disorders, diabetes, dislipidemia, colonic hypersensitivity associated with psychopathological disturbance, and stress. It further relates to the use of such compounds in treating physiological disorders induced or facilitated by CRF or those associated with an excess of neuropeptide Y. In particular, this invention provides aminoalkyl substituted 9H-pyridino[2,3-b]indole and 9H-pyrimidino[4,5-b]indole derivatives of Formula I which selectively bind to corticotropin-releasing factor (CRF₁) receptors and/or to mammalian neuropeptide Y (NPY₁) receptors.

Accordingly, a broad embodiment of the invention is directed to a compound of Formula I:



wherein


Ar is phenyl, 1- or 2-naphthyl, 2-, 3-, or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, optionally mono-, di-, or tri-substituted with halogen, trifluoromethyl, hydroxy, amino, lower alkylamino, lower dialkylamino, carboxamido, N-lower alkyl

carboxamido, N,N-lower dialkyl carboxamido, C₁-C₆ alkyl, C₁-C₆ alkoxy, with the proviso that at least one of the positions ortho or para to the point of attachment of Ar to the tricyclic ring system is substituted;

5 R¹ is hydrogen, halogen, trifluoromethyl, C₁-C₆ alkyl, or (C₁-C₆ alkyl)-G¹-R² where G¹ is oxygen or sulfur and R² is hydrogen or C₁-C₆ alkyl;

W is N or C-R³ where R³ is hydrogen or C₁-C₆ alkyl; and

X is



10

(i)

wherein

V¹ and V² are CH₂, CO, CS, SO₂ or CH(C₁-C₆ alkyl), with the proviso that both V¹ and V² cannot both be CO, CS or SO₂;

Y¹ and Y² independently represent a bond or C₁-C₆ alkylene;

15

A¹ is NR⁴R⁵ wherein R⁴ and R⁵ are independently hydrogen, a lower alkyl group which optionally forms a heterocycloalkyl group with Y¹;


C₁-C₆ arylalkyl or C₁-C₆ heteroarylalkyl, where aryl is phenyl, and heteroaryl is 2-, 3-, or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or 1-, 2- or 5-tetrazolyl, each of which is optionally mono-, di-, or tri-substituted with halogen, trifluoromethyl, hydroxy, amino, lower alkylamino, lower dialkylamino, lower

25

alkylcarboxamido, lower alkyl, lower alkoxy, with the proviso that 2 adjacent substituents can together form a 5-7 fused cycloalkyl or heterocycloalkyl ring;

lower alkanoyl, lower alkylsulfonyl, with the proviso that R^4 and R^5 cannot both be alkanoyl or alkylsulfonyl; or

NR^4R^5 taken together form a C₃-C₆ heterocycloalkyl or a group of the formula:



wherein e and f are independently 1, 2 or 3 and the sum of e and f is at least 3; and

G^2 is

NR^6 wherein R^6 is hydrogen or C₁-C₆ alkyl, or

$CH(C_0-C_6 \text{ alkylene})-G^3-R^7$ wherein G^3 is CONH,


CONH(C₁-C₆ alkyl), NH, NH(C₁-C₆ alkyl) and R^7 is hydrogen or C₁-C₆ alkyl; or

CONH₂, CO[N(C₁-C₆ alkyl) R^8] wherein R^8 is hydrogen or C₁-C₆ alkyl;

C₁-C₆ arylalkyl or C₁-C₆ heteroarylalkyl, where aryl is phenyl, and heteroaryl is 2-, 3-, or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or 1-, 2- or 5-tetrazolyl, each of which is optionally mono-, di-, or tri-substituted with halogen,

trifluoromethyl, hydroxy, amino, lower
alkylamino, lower dialkylamino, lower
alkylcarboxamido, lower alkyl, lower alkoxy,
with the proviso that 2 adjacent substituents
can together form a 5-7 fused cycloalkyl or
heterocycloalkyl ring;

A^2 is hydrogen, C_1 - C_6 alkyl, (C_1 - C_6 alkylene)- G^4 - R^9
wherein G^4 is oxygen or sulfur and R^9 is hydrogen,
trifluoromethyl or C_1 - C_6 alkyl;




(ii)

wherein heteroaryl is 2-, 3- or 4-pyridyl, 2-, 4- or 5-
pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-
oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-
pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or 1-,
2- or 5-tetrazolyl, each of which is optionally mono-
or disubstituted with halogen, trifluoromethyl,
amino, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, with the proviso
that tetrazolyl can have at most one substituent;

Z^1 is C_1 - C_6 alkyl; and

V^2 , Y^2 and A^2 are as defined above;



(iii)

where

Z^2 is carbon or nitrogen;

where

when Z^2 is CH, n is 0, 1, 2 or 3 and p is 1, 2, or 3,

R^{10} is carboxamido, or (C₀-C₆ alkylene)-G⁵-R¹¹

wherein G⁵ is NH, NH(C₁-C₆ alkyl) and R¹¹ is hydrogen, C₁-C₆ alkyl, C₁-C₆ arylalkyl or C₁-C₆ heteroarylalkyl, where aryl is phenyl, and heteroaryl is 2-, 3-, or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or 1-, 2- or 5-tetrazolyl, each of which is optionally mono-, di-, or tri-substituted with halogen, trifluoromethyl, hydroxy, amino, lower alkylamino, lower dialkylamino, lower alkylcarboxamido, lower alkyl, lower alkoxy, with the proviso that 2 adjacent substituents can together form a 5-7 fused cycloalkyl or heterocycloalkyl ring; ;

when Z^2 is carbon, n is 1 or 2 and p is 1 or 2, R¹⁰ is amino; or

when Z^2 is nitrogen, n is 1 or 2 and p is 1 or 2, R¹⁰ is hydrogen; or

(iv) a nitrogen heterocycle of the formula:



wherein the N-ring represents triazolyl, tetrazolyl, imidazolyl, or pyrazolyl, each of which is optionally substituted with amino, trifluoromethyl, carboxamido, or (C₁-C₆

alkylene)-G⁶-R¹² wherein G⁶ is NH, NH(C₁-C₆ alkyl) and R¹² is hydrogen, C₁-C₆ alkyl, C₁-C₆ arylalkyl or C₁-C₆ heteroarylalkyl, where aryl is phenyl, and heteroaryl is 2-, 3-, or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or 1-, 2- or 5-tetrazolyl, each of which is optionally mono-, di-, or tri-substituted with halogen, trifluoromethyl, hydroxy, amino, lower alkylamino, lower dialkylamino, lower alkylcarboxamido, lower alkyl, lower alkoxy, with the proviso that 2 adjacent substituents can together form a 5-7 fused cycloalkyl or heterocycloalkyl ring.

The compounds of Formula I are antagonists at the CRF₁ receptor and are useful in the diagnosis and treatment of stress related disorders such as post traumatic stress disorder (PTSD) as well as depression, headache and anxiety.


They are useful in methods for the treatment of psychiatric and affective disorders and neurological diseases, including major depression, headaches, anxiety-related disorders, post-traumatic stress disorder, supranuclear palsy, as well as treatment of immunological, cardiovascular or heart-related diseases, hypertension, feeding disorders, diabetes, dislipidemia, colonic hypersensitivity associated with psychopathological disturbance, and stress. Such methods involve administration to a mammal of an effective amount of a compound of the invention.

The compounds of Formula I are also neuropeptide Y_1 receptor antagonists, and, therefore, are also of value in the treatment of a wide variety of clinical conditions which are characterized by the presence of an excess of neuropeptide Y.

5 The compounds of Formula I have different chemical structures which affect their selectivity towards either the CRF_1 or the NPY_1 receptors.

DETAILED DESCRIPTION OF THE INVENTION

The novel compounds encompassed by the instant invention can be described by general Formula I:




wherein

Ar is phenyl, 1- or 2-naphthyl, 2-, 3-, or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, optionally mono-, di-, or tri-substituted with halogen, trifluoromethyl, hydroxy, amino, lower alkylamino, lower dialkylamino, carboxamido, N-lower alkyl carboxamido, N,N-lower dialkyl carboxamido, C₁-C₆ alkyl, C₁-C₆ alkoxy, with the proviso that at least one of the positions ortho or para to the point of attachment of Ar to the tricyclic ring system is substituted;

R¹ is hydrogen, halogen, trifluoromethyl, C₁-C₆ alkyl, or (C₁-C₆ alkyl)-G¹-R² where G¹ is oxygen or sulfur and R² is hydrogen or C₁-C₆ alkyl;

W is N or C-R³ where R³ is hydrogen or C₁-C₆ alkyl; and

X is



(i)

wherein

V¹ and V² are CH₂, CO, CS, SO₂ or CH(C₁-C₆ alkyl), with the proviso that both V¹ and V² cannot both be CO, CS or SO₂;


Y¹ and Y² independently represent a bond or C₁-C₆ alkylene;

A¹ is NR⁴R⁵ wherein R⁴ and R⁵ are independently hydrogen, a lower alkyl group which optionally forms a heterocycloalkyl group with Y¹;

C₁-C₆ arylalkyl or C₁-C₆ heteroarylalkyl, where aryl is phenyl, and heteroaryl is 2-, 3-, or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or 1-, 2- or 5-tetrazolyl, each of which is optionally mono-, di-, or tri-substituted with halogen, trifluoromethyl, hydroxy, amino, lower alkylamino, lower dialkylamino, lower alkylcarboxamido, lower alkyl, lower alkoxy, with the proviso that 2 adjacent substituents can together form a 5-7 fused cycloalkyl or heterocycloalkyl ring;

lower alkanoyl, lower alkylsulfonyl, with the proviso that R⁴ and R⁵ cannot both be alkanoyl or alkylsulfonyl; or

NR⁴R⁵ taken together form a C₃-C₆ heterocycloalkyl or a group of the formula:



wherein e and f are independently 1, 2 or 3 and the sum of e and f is at least 3; and

G² is


NR⁶ wherein R⁶ is hydrogen or C₁-C₆ alkyl, or

CH(C₀-C₆ alkylene)-G³-R⁷ wherein G³ is CONH,
 CONH(C₁-C₆ alkyl), NH, NH(C₁-C₆ alkyl) and R⁷
 is hydrogen or C₁-C₆ alkyl; or

CONH₂, CO[N(C₁-C₆ alkyl)R⁸] wherein R⁸ is
 hydrogen or C₁-C₆ alkyl;

C₁-C₆ arylalkyl or C₁-C₆ heteroarylalkyl, where
 aryl is phenyl, and heteroaryl is 2-, 3-, or 4-
 pyridyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 4-
 imidazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-
 thiazolyl, 1-, 3- or 4-pyrazolyl, 1-, 3- or 4-
 triazolyl, 2-pyrazinyl, or 1-, 2- or 5-
 tetrazolyl, each of which is optionally mono-,
 di-, or tri-substituted with halogen,
 trifluoromethyl, hydroxy, amino, lower
 alkylamino, lower dialkylamino, lower
 alkylcarboxamido, lower alkyl, lower alkoxy,
 with the proviso that 2 adjacent substituents
 can together form a 5-7 fused cycloalkyl or
 heterocycloalkyl ring;

A² is hydrogen, C₁-C₆ alkyl, (C₁-C₆ alkylene)-G⁴-R⁹
 wherein G⁴ is oxygen or sulfur and R⁹ is hydrogen,
 trifluoromethyl or C₁-C₆ alkyl;




(ii)

wherein heteroaryl is 2-, 3- or 4-pyridyl, 2-, 4- or 5-
 pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-
 oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-
 pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or 1-,

2- or 5-tetrazolyl, each of which is optionally mono- or disubstituted with halogen, trifluoromethyl, amino, C₁-C₆ alkyl, C₁-C₆ alkoxy, with the proviso that tetrazolyl can have at most one substituent;

5 Z¹ is C₁-C₆ alkyl; and

V², Y² and A² are as defined above;



(iii)

where

Z² is carbon or nitrogen;

10 where

when Z² is CH, n is 0, 1, 2 or 3 and p is 1, 2, or 3,

R¹⁰ is carboxamido, or (C₀-C₆ alkylene)-G⁵-R¹¹

wherein G⁵ is NH, NH(C₁-C₆ alkyl) and R¹¹ is

hydrogen, C₁-C₆ alkyl, C₁-C₆ arylalkyl or C₁-C₆

15 heteroarylalkyl, where aryl is phenyl, and

heteroaryl is 2-, 3-, or 4-pyridyl, 2-, 4- or 5-

pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-

oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-

pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or

20 1-, 2- or 5-tetrazolyl, each of which is

optionally mono-, di-, or tri-substituted with

halogen, trifluoromethyl, hydroxy, amino, lower

alkylamino, lower dialkylamino, lower

alkylcarboxamido, lower alkyl, lower alkoxy, with

25 the proviso that 2 adjacent substituents can

together form a 5-7 fused cycloalkyl or heterocycloalkyl ring; ;

when Z^2 is carbon, n is 1 or 2 and p is 1 or 2, R^{10} is amino; or


5 when Z^2 is nitrogen, n is 1 or 2 and p is 1 or 2, R^{10} is hydrogen; or

(iv) a nitrogen heterocycle of the formula:



wherein the N-ring represents triazolyl, tetrazolyl, imidazolyl, or pyrazolyl, each of which is optionally substituted with amino, trifluoromethyl, carboxamido, or (C₁-C₆ alkylene)-G⁶-R¹² wherein G⁶ is NH, NH(C₁-C₆ alkyl) and R¹² is hydrogen, C₁-C₆ alkyl, C₁-C₆ arylalkyl or C₁-C₆ heteroarylalkyl, where aryl is phenyl, and heteroaryl is 2-, 3-, or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or 1-, 2- or 5-tetrazolyl, each of which is optionally mono-, di-, or tri-substituted with halogen, trifluoromethyl, hydroxy, amino, lower alkylamino, lower dialkylamino, lower alkylcarboxamido, lower alkyl, lower alkoxy, with the proviso that 2 adjacent substituents can together form a 5-7 fused cycloalkyl or heterocycloalkyl ring.

Preferred compounds of Formula I include those of Formula IA:




IA

- 5 wherein
 each R_a is C_1 - C_6 alkyl;
 R_b is hydrogen or methyl;
 R_1 is C_1 - C_6 alkyl;
 R_s is C_1 - C_6 alkyl, $(C_3$ - $C_5)$ cycloalkyl $(C_1$ - $C_3)$ alkyl, $(C_1$ - $C_3)$ alkoxy $(C_1$ -
 10 $C_3)$ alkyl, or $(C_3$ - $C_5)$ cycloalkyl;
 t is 1, 2 or 3; and
 R_x is hydrogen, C_1 - C_6 alkyl, phenyl $(C_1$ - $C_6)$ alkyl where phenyl is
 optionally mono- or disubstituted independently with C_1 - C_6
 alkyl, C_1 - C_6 alkoxy, halogen, or hydroxy; and
 15 R_y is hydrogen, C_1 - C_6 alkyl, $(C_3$ - $C_6)$ cycloalkyl; or
 NR_xR_y represents pyrrolidinyl, N - $(C_1$ - $C_6)$ alkylpyrrolidin-2-yl,
 piperidinyl, morpholinyl, or N - $(C_1$ - $C_6)$ alkylpiperazinyl.

Preferred compounds of Formula IA include those where R_s
 20 is C_1 - C_6 alkyl or cyclopropylmethyl. Other preferred compounds
 of Formula IA include those where R_s is cyclopropyl $(C_1$ - $C_3)$ alkyl.
 Still other preferred compounds of Formula IA include those
 where R_x and R_y independently represent hydrogen or C_1 - C_2 alkyl.
 More preferred compounds of IA include those where R_s is
 25 cyclopropyl $(C_1$ - $C_3)$ alkyl, R_x and R_y independently represent
 hydrogen or C_1 - C_2 alkyl; and each R_a is methyl. Particularly
 preferred compounds of IA are those W is nitrogen.

Other preferred compounds of Formula I include those of

Formula IB:




IB

wherein

- 5 each R_a is C_1 - C_6 alkyl;
 R_1 is C_1 - C_6 alkyl;
 R_s is C_1 - C_6 alkyl, cyclopropyl(C_1 - C_3)alkyl or (C_1 - C_3)alkoxy(C_1 - C_3)alkyl;
 t is 1 or 2;
 10 R_x is hydrogen, C_1 - C_6 alkyl, phenyl(C_1 - C_6)alkyl where phenyl is optionally mono- or disubstituted independently with C_1 - C_6 alkyl, C_1 - C_6 alkoxy, halogen, or hydroxy; and
 R_y is hydrogen, C_1 - C_6 alkyl, (C_3 - C_6)cycloalkyl; or
 NR_xR_y represents pyrrolidinyl, N-(C_1 - C_6)alkylpyrrolidin-2-yl,
 15 piperidinyl, morpholinyl, or N-(C_1 - C_6)alkylpiperazinyl.

- Preferred compounds of Formula IB include those where R_s is C_1 - C_6 alkyl or cyclopropylmethyl. Other preferred compounds of Formula IB include those where R_s is cyclopropyl(C_1 - C_3)alkyl.
 20 Yet other preferred compounds of Formula IB include those where t is 1 and R_x and R_y independently represent hydrogen or C_1 - C_2 alkyl. More preferred compounds of Formula IB include those where R_s is cyclopropyl(C_1 - C_3)alkyl, t is 1 and R_x and R_y independently represent hydrogen or C_1 - C_2 alkyl. Particularly
 25 preferred compounds of IB are those where each R_a is methyl, R_s is cyclopropyl(C_1 - C_3)alkyl, t is 1 and R_x and R_y independently represent hydrogen or C_1 - C_2 alkyl. Highly preferred compounds of Formula IB are those where W is nitrogen.

Other preferred compounds of Formula I include those of Formula IC:



IC

5 wherein

each R_a is C_1 - C_6 alkyl;


R_1 is C_1 - C_6 alkyl;

t is 1 or 2;

R_x and R_y are different and represent hydrogen; (C_3 -
 10 C_7)cycloalkylamino(C_1 - C_3)alkyl, carboxamido, (C_3 -
 C_7)cycloalkylamino, C_2 - C_6 alkanoyl optionally substituted
 in the ω -position with C_1 - C_6 alkyl or phenyl optionally
 mono- or disubstituted independently with C_1 - C_6 alkyl, C_1 -
 C_6 alkoxy, halogen, or hydroxy, provided that at least one
 15 of R_x and R_y is hydrogen.

Preferred compounds of Formula IC include those wherein R^1
 is C_1 - C_2 alkyl and W is nitrogen. Other preferred compounds of
 IC are those where each R_a is methyl and W is nitrogen.

20 Other preferred compounds of the invention have Formula II




II

wherein

each R_a independently represents C_1-C_6 alkyl;

R^1 is hydrogen, halogen, trifluoromethyl, C_1-C_6 alkyl, or $(C_1-C_6 \text{ alkyl})-G^1-R^2$ where G^1 is oxygen or sulfur and R^2 is hydrogen or C_1-C_6 alkyl;

X is



(i)

wherein


V^1 and V^2 are CH_2 , CO, CS, SO_2 or $CH(C_1-C_6 \text{ alkyl})$, with the proviso that both V^1 and V^2 cannot both be CO, CS or SO_2 ;

Y^1 and Y^2 independently represent a bond or C_1-C_6 alkylene;

A^1 is NR^4R^5 wherein R^4 and R^5 are independently hydrogen, a lower alkyl group which optionally forms a heterocycloalkyl group with Y^1 ;

C_1-C_6 arylalkyl or C_1-C_6 heteroarylalkyl, where aryl is phenyl, and heteroaryl is 2-, 3-, or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or 1-, 2- or 5-tetrazolyl, each of which is optionally mono-, di-, or tri-substituted with halogen, trifluoromethyl, hydroxy, amino, lower alkylamino, lower dialkylamino, lower alkylcarboxamido, lower alkyl, lower alkoxy, with the proviso that 2 adjacent substituents

can together form a 5-7 fused cycloalkyl or heterocycloalkyl ring;
 lower alkanoyl, lower alkylsulfonyl, with the proviso
 that R^4 and R^5 cannot both be alkanoyl or
 alkylsulfonyl; or
 NR^4R^5 taken together form a C₃-C₆ heterocycloalkyl or
 a group of the formula:



wherein e and f are independently 1, 2 or 3 and
 the sum of e and f is at least 3; and

G^2 is

NR^6 wherein R^6 is hydrogen or C₁-C₆ alkyl, or

$CH(C_0-C_6 \text{ alkylene})-G^3-R^7$ wherein G^3 is CONH,


CONH(C₁-C₆ alkyl), NH, NH(C₁-C₆ alkyl) and R^7
 is hydrogen or C₁-C₆ alkyl; or

CONH₂, CO[N(C₁-C₆ alkyl) R^8] wherein R^8 is
 hydrogen or C₁-C₆ alkyl;

C₁-C₆ arylalkyl or C₁-C₆ heteroarylalkyl, where
 aryl is phenyl, and heteroaryl is 2-, 3-, or 4-
 pyridyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 4-
 imidazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-
 thiazolyl, 1-, 3- or 4-pyrazolyl, 1-, 3- or 4-
 triazolyl, 2-pyrazinyl, or 1-, 2- or 5-
 tetrazolyl, each of which is optionally mono-,
 di-, or tri-substituted with halogen,
 trifluoromethyl, hydroxy, amino, lower
 alkylamino, lower dialkylamino, lower

alkylcarboxamido, lower alkyl, lower alkoxy, with the proviso that 2 adjacent substituents can together form a 5-7 fused cycloalkyl or heterocycloalkyl ring;


- 5 A^2 is hydrogen, C_1-C_6 alkyl, $(C_1-C_6 \text{ alkylene})-G^4-R^9$ wherein G^4 is oxygen or sulfur and R^9 is hydrogen, trifluoromethyl or C_1-C_6 alkyl;



- 10 wherein heteroaryl is 2-, 3- or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or 1-, 2- or 5-tetrazolyl, each of which is optionally mono-
15 or disubstituted with halogen, trifluoromethyl, amino, C_1-C_6 alkyl, C_1-C_6 alkoxy, with the proviso that tetrazolyl can have at most one substituent;

Z^1 is C_1-C_6 alkyl; and

V^2 , Y^2 and A^2 are as defined above;



- 20 (iii)
where

Z^2 is carbon or nitrogen;

where

when Z^2 is CH, n is 0, 1, 2 or 3 and p is 1, 2, or 3,
 R^{10} is carboxamido, or $(C_0-C_6 \text{ alkylene})-G^5-R^{11}$
 wherein G^5 is NH, $NH(C_1-C_6 \text{ alkyl})$ and R^{11} is
 hydrogen, C_1-C_6 alkyl, C_1-C_6 arylalkyl or C_1-C_6
 heteroarylalkyl, where aryl is phenyl, and
 heteroaryl is 2-, 3-, or 4-pyridyl, 2-, 4- or 5-
 pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-
 oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-
 pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or
 1-, 2- or 5-tetrazolyl, each of which is
 optionally mono-, di-, or tri-substituted with
 halogen, trifluoromethyl, hydroxy, amino, lower
 alkylamino, lower dialkylamino, lower
 alkylcarboxamido, lower alkyl, lower alkoxy, with
 the proviso that 2 adjacent substituents can
 together form a 5-7 fused cycloalkyl or
 heterocycloalkyl ring; ;

when Z^2 is carbon, n is 1 or 2 and p is 1 or 2, R^{10} is
 amino; or

when Z^2 is nitrogen, n is 1 or 2 and p is 1 or 2, R^{10} is
 hydrogen; or

(iv) a nitrogen heterocycle of the formula:




wherein the N-ring represents triazolyl, tetrazolyl,
 imidazolyl, or pyrazolyl, each of which is
 optionally substituted with amino,
 trifluoromethyl, carboxamido, or $(C_1-C_6$
 alkylene)- G^6-R^{12} wherein G^6 is NH, $NH(C_1-C_6$
 alkyl) and R^{12} is hydrogen, C_1-C_6 alkyl, C_1-C_6

arylalkyl or C₁-C₆ heteroarylalkyl, where aryl is phenyl, and heteroaryl is 2-, 3-, or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or 1-, 2- or 5-tetrazolyl, each of which is optionally mono-, di-, or tri-substituted with halogen, trifluoromethyl, hydroxy, amino, lower alkylamino, lower dialkylamino, lower alkylcarboxamido, lower alkyl, lower alkoxy, with the proviso that 2 adjacent substituents can together form a 5-7 fused cycloalkyl or heterocycloalkyl ring.

Preferred compounds of Formula II are those where V¹ and V² represent methylene; Y¹ is a bond; A¹ represents pyrrolidinyl, morpholinyl, piperazinyl, mono- or di-C₁-C₆ alkyl, C₁-C₆ arylalkyl or C₁-C₆ heteroarylalkyl, where aryl is phenyl, and heteroaryl is 2-, 3-, or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or 1-, 2- or 5-tetrazolyl, each of which is optionally mono-, di-, or tri-substituted with halogen, trifluoromethyl, hydroxy, amino, lower alkylamino, lower dialkylamino, lower alkylcarboxamido, lower alkyl, lower alkoxy, with the proviso that 2 adjacent substituents can together form a 5-7 fused cycloalkyl or heterocycloalkyl ring; Y² represents a bond or methylene; and A² represents C₁-C₆ alkyl or C₁-C₆ alkoxymethyl.

Other preferred compounds of the invention have Formula III




III

wherein

each R_a independently represents C_1 - C_6 alkyl;

5 R^1 is hydrogen, halogen, trifluoromethyl, C_1 - C_6 alkyl, or $(C_1$ - C_6 alkyl)- G^1 - R^2 where G^1 is oxygen or sulfur and R^2 is hydrogen or C_1 - C_6 alkyl; and

X is



(i)

wherein

10 V^1 and V^2 are CH_2 , CO, CS, SO_2 or $CH(C_1$ - C_6 alkyl), with the proviso that both V^1 and V^2 cannot both be CO, CS or SO_2 ;

Y^1 and Y^2 independently represent a bond or C_1 - C_6 alkylene;


15 A^1 is NR^4R^5 wherein R^4 and R^5 are independently hydrogen, a lower alkyl group which optionally forms a heterocycloalkyl group with Y^1 ;

20 C_1 - C_6 arylalkyl or C_1 - C_6 heteroarylalkyl, where aryl is phenyl, and heteroaryl is 2-, 3-, or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-pyrazolyl, 1-, 3- or 4-

triazolyl, 2-pyrazinyl, or 1-, 2- or 5-tetrazolyl, each of which is optionally mono-, di-, or tri-substituted with halogen, trifluoromethyl, hydroxy, amino, lower alkylamino, lower dialkylamino, lower alkylcarboxamido, lower alkyl, lower alkoxy, with the proviso that 2 adjacent substituents can together form a 5-7 fused cycloalkyl or heterocycloalkyl ring;

lower alkanoyl, lower alkylsulfonyl, with the proviso that R^4 and R^5 cannot both be alkanoyl or alkylsulfonyl; or

NR^4R^5 taken together form a C_3 - C_6 heterocycloalkyl or a group of the formula:



wherein e and f are independently 1, 2 or 3 and the sum of e and f is at least 3; and

G^2 is

NR^6 wherein R^6 is hydrogen or C_1 - C_6 alkyl, or


$CH(C_0$ - C_6 alkylene)- G^3 - R^7 wherein G^3 is CONH, CONH(C_1 - C_6 alkyl), NH, NH(C_1 - C_6 alkyl) and R^7 is hydrogen or C_1 - C_6 alkyl; or

CONH₂, CO[N(C_1 - C_6 alkyl) R^8] wherein R^8 is hydrogen or C_1 - C_6 alkyl;

C_1 - C_6 arylalkyl or C_1 - C_6 heteroarylalkyl, where aryl is phenyl, and heteroaryl is 2-, 3-, or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 4-

imidazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or 1-, 2- or 5-tetrazolyl, each of which is optionally mono-, di-, or tri-substituted with halogen, trifluoromethyl, hydroxy, amino, lower alkylamino, lower dialkylamino, lower alkylcarboxamido, lower alkyl, lower alkoxy, with the proviso that 2 adjacent substituents can together form a 5-7 fused cycloalkyl or heterocycloalkyl ring;


A^2 is hydrogen, C_1 - C_6 alkyl, (C_1 - C_6 alkylene)- G^4 - R^9 wherein G^4 is oxygen or sulfur and R^9 is hydrogen, trifluoromethyl or C_1 - C_6 alkyl;



wherein heteroaryl is 2-, 3- or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or 1-, 2- or 5-tetrazolyl, each of which is optionally mono- or disubstituted with halogen, trifluoromethyl, amino, C_1 - C_6 alkyl, C_1 - C_6 alkoxy, with the proviso that tetrazolyl can have at most one substituent;

Z^1 is C_1 - C_6 alkyl; and

V^2 , Y^2 and A^2 are as defined above;



(iii)

where

 Z^2 is carbon or nitrogen;

where

5 when Z^2 is CH, n is 0, 1, 2 or 3 and p is 1, 2, or 3,

R^{10} is carboxamido, or (C₀-C₆ alkylene)-G⁵-R¹¹

wherein G⁵ is NH, NH(C₁-C₆ alkyl) and R¹¹ is
 10 hydrogen, C₁-C₆ alkyl, C₁-C₆ arylalkyl or C₁-C₆
 heteroarylalkyl, where aryl is phenyl, and
 heteroaryl is 2-, 3-, or 4-pyridyl, 2-, 4- or 5-
 pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-
 oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-
 pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or
 1-, 2- or 5-tetrazolyl, each of which is
 15 optionally mono-, di-, or tri-substituted with
 halogen, trifluoromethyl, hydroxy, amino, lower
 alkylamino, lower dialkylamino, lower
 alkylcarboxamido, lower alkyl, lower alkoxy, with
 the proviso that 2 adjacent substituents can
 20 together form a 5-7 fused cycloalkyl or
 heterocycloalkyl ring; ;

when Z^2 is carbon, n is 1 or 2 and p is 1 or 2, R^{10} is
 amino; or

when Z^2 is nitrogen, n is 1 or 2 and p is 1 or 2, R^{10} is
 25 hydrogen; or

(iv) a nitrogen heterocycle of the formula:



wherein the N-ring represents triazolyl, tetrazolyl, imidazolyl, or pyrazolyl, each of which is optionally substituted with amino, trifluoromethyl, carboxamido, or (C₁-C₆ alkylene)-G⁶-R¹² wherein G⁶ is NH, NH(C₁-C₆ alkyl) and R¹² is hydrogen, C₁-C₆ alkyl, C₁-C₆ arylalkyl or C₁-C₆ heteroarylalkyl, where aryl is phenyl, and heteroaryl is 2-, 3-, or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or 1-, 2- or 5-tetrazolyl, each of which is optionally mono-, di-, or tri-substituted with halogen, trifluoromethyl, hydroxy, amino, lower alkylamino, lower dialkylamino, lower alkylcarboxamido, lower alkyl, lower alkoxy, with the proviso that 2 adjacent substituents can together form a 5-7 fused cycloalkyl or heterocycloalkyl ring.

Preferred compounds of Formula III are those where V¹ and V² represent methylene; Y¹ is a bond; A¹ represents pyrrolidinyl, morpholinyl; piperazinyl, mono- or di-C₁-C₆ alkyl, C₁-C₆ arylalkyl or C₁-C₆ heteroarylalkyl, where aryl is phenyl, and heteroaryl is 2-, 3-, or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or 1-, 2- or 5-tetrazolyl, each of

which is optionally mono-, di-, or tri-substituted with halogen, trifluoromethyl, hydroxy, amino, lower alkylamino, lower dialkylamino, lower alkylcarboxamido, lower alkyl, lower alkoxy, with the proviso that 2 adjacent substituents can together form a 5-7 fused cycloalkyl or heterocycloalkyl ring; Y² represents a bond or methylene; and A² represents C₁-C₆ alkyl or C₁-C₆ alkoxyethyl.

Preferred compounds of the invention include:

10 4-(N-(2-N',N'-Dimethylaminoethyl)-N-ethyl)amino-2-methyl-9-(4-bromo-2,6-dimethylphenyl)-9H-pyrimidino[4,5-b]indole

4-(N-(2-N',N'-Dimethylaminoethyl)-N-propyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole

15

4-(N-(2-N',N'-Dimethylaminoethyl)-N-2-methoxyethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole

20

4-(N-(2-N',N'-Dimethylaminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole

25

4-(N-(2-Pyrrolidinoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole

4-(N-(2-N'-Ethylaminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole

30

4-(N-(2-N'-Ethyl-N'-methylaminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole

4 - (N - (2 - N' , N' - Dimethylaminoethyl) - N -
cyclopropylmethyl) amino - 2 - methyl - 9 - (4 - methoxy - 2 - methylphenyl) -
9H - pyridino [2, 3 - b] indole

5 4 - (N - (2 - N' , N' - Diethylaminoethyl) - N -
cyclopropylmethyl) amino - 2 - methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H -
pyridino [2, 3 - b] indole

10 4 - (N - (2 - N' - Methylaminoethyl) - N - cyclopropylmethyl) amino - 2 -
methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H - pyridino [2, 3 - b] indole

4 - (N - (2 - Aminoethyl) - N - cyclopropylmethyl) amino - 2 - methyl - 9 -
(2, 4, 6 - trimethylphenyl) - 9H - pyridino [2, 3 - b] indole

15 4 - (N - (2 - Piperidinoethyl) - N - cyclopropylmethyl) amino - 2 -
methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H - pyridino [2, 3 - b] indole

20 4 - (N - (2 - Pyrrolidinoethyl) - N - propyl) amino - 2 - methyl - 9 -
(2, 4, 6 - trimethylphenyl) - 9H - pyridino [2, 3 - b] indole

4 - (N - (2 - Piperidinoethyl) - N - propyl) amino - 2 - methyl - 9 -
(2, 4, 6 - trimethylphenyl) - 9H - pyridino [2, 3 - b] indole

25 4 - (N - (2 - Morpholinoethyl) - N - cyclopropylmethyl) amino - 2 -
methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H - pyridino [2, 3 - b] indole

4 - (N - (2 - N' - Ethyl - N' - methylaminoethyl) - N - propyl) amino - 2 -
methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H - pyridino [2, 3 - b] indole

30 4 - (N - (2 - N' , N' - Dimethylaminoethyl) - N -
cyclopropylmethyl) amino - 2 - methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H -
pyrimidino [4, 5 - b] indole

4 - (N - (2 - (1-Imidazolyl) ethyl) -N-cyclopropylmethyl) amino-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole

5 4 - (N - (2-Pyrrolidinoethyl) -N-1-oxopropyl) amino-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole

4 - (N - (2-Pyrrolidinoethyl) -N-cyclopropyloxomethyl) amino-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole

10 4 - (N-2 - (N' -Methylpiperazinyl) ethyl-N-cyclopropylmethyl) - amino-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole

15 4 - (N - (2-Pyridylmethyl) -N-cyclopropylmethyl) amino-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole

4 - Piperazinyl-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole

20 4 - (N - (2-Aminoethyl) -N-propyl) amino-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole

4 - (N - (2-Aminoethyl) -N-ethyl) amino-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole

25

4 - (N - (2 - (4-Triazolyl) ethyl) -N-cyclopropylmethyl) amino-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole

30 4 - (N - (2-Pyrrolidinoethyl) -N-cyclopropylmethyl) amino-2-methyl-9 - (2,4-dimethylphenyl) -9H-pyridino[2,3-b] indole

4 - (N - (3-Pyrrolidinoproyl) -N-cyclopropylmethyl) amino-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole

4 - (N - (2 - (2 - Phenethylamino) ethyl) - N -
cyclopropylmethyl) amino - 2 - methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H -
pyridino [2, 3 - b] indole

5 4 - (N - (2 - Pyrrolidinoethyl) - N - cyclopropylmethyl) amino - 2 -
methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H - pyrimidino [4, 5 - b] indole

4 - (N - (2 - Pyrrolidinoethyl) - N - 2 - methoxyethyl) amino - 2 -
methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H - pyridino [2, 3 - b] indole

10

4 - (N - (2 - Pyrrolidinoethyl) - N - ethyl) amino - 2 - methyl - 9 -
(2, 4, 6 - trimethylphenyl) - 9H - pyridino [2, 3 - b] indole

4 - (N - (2 - Pyrrolidinoethyl) - N - butyl) amino - 2 - methyl - 9 -
15 (2, 4, 6 - trimethylphenyl) - 9H - pyridino [2, 3 - b] indole

4 - (N - (2 - Pyrrolidinoethyl) - N - 2 - methylpropyl) amino - 2 -
methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H - pyridino [2, 3 - b] indole

20 4 - (N - (2 - Carboxamidoethyl) - N - cyclopropylmethyl) amino - 2 -
methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H - pyridino [2, 3 - b] indole

4 - (N - (2 - Pyrrolidinoethyl) - N - cyclopropylmethyl) amino - 2 -
methyl - 9 - (4 - bromo - 2, 6 - dimethylphenyl) - 9H - pyrimidino [4, 5 -
25 b] indole

4 - (N - (2 - N' , N' - Dimethylaminoethyl) - N - 2 - methylpropyl) amino -
2 - methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H - pyridino [2, 3 - b] indole

30 4 - (N - (2 - N' - Ethyl - N' - methylaminoethyl) - N - 2 -
methylpropyl) amino - 2 - methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H -
pyridino [2, 3 - b] indole

4 - (N - (2 - N' , N' - Dimethylaminoethyl) - N - propyl) amino - 2 -
methyl - 9 - (2 , 4 , 6 - trimethylphenyl) - 9H - pyrimidino [4 , 5 - b] indole

4 - (N - (2 - N' , N' - Dimethylaminoethyl) - N - butyl) amino - 2 -
5 methyl - 9 - (2 , 4 , 6 - trimethylphenyl) - 9H - pyrimidino [4 , 5 - b] indole

4 - (N - (2 - N' , N' - Dimethylaminoethyl) - N - cyclopropyl) amino - 2 -
methyl - 9 - (2 , 4 , 6 - trimethylphenyl) - 9H - pyrimidino [4 , 5 - b] indole

10 4 - (N - (2 - (1 - Methyl - 2 - pyrrolidino) ethyl) - N -
cyclopropylmethyl) amino - 2 - methyl - 9 - (2 , 4 , 6 - trimethylphenyl) - 9H -
pyridino [2 , 3 - b] indole

4 - (N - (2 - N' , N' - Dimethylaminoethyl) - N - cyclobutyl) amino - 2 -
15 methyl - 9 - (2 , 4 , 6 - trimethylphenyl) - 9H - pyrimidino [4 , 5 - b] indole

4 - (N - (2 - N' , N' - Dimethylaminoethyl) - N - isopropyl) amino - 2 -
methyl - 9 - (2 , 4 , 6 - trimethylphenyl) - 9H - pyrimidino [4 , 5 - b] indole

20 4 - (N - (2 - Pyrrolidinoethyl) - N - isopropyl) amino - 2 - methyl - 9 -
(2 , 4 , 6 - trimethylphenyl) - 9H - pyrimidino [4 , 5 - b] indole

4 - (N - (2 - N' , N' - Dimethylaminoethyl) - N - isopropyl) amino - 2 -
methyl - 9 - (2 , 4 , 6 - trimethylphenyl) - 9H - pyrimidino [4 , 5 - b] indole

25

In certain situations, the compounds of Formula I may
contain one or more asymmetric carbon atoms, so that the
compounds can exist in different stereoisomeric forms. These
compounds can be, for example, racemates or optically active
30 forms. In these situations, the single enantiomers, i.e.,
optically active forms, can be obtained by asymmetric synthesis
or by resolution of the racemates. Resolution of the racemates
can be accomplished, for example, by conventional methods such

as crystallization in the presence of a resolving agent, or chromatography, using, for example a chiral HPLC column.

Representative compounds of the present invention, which are encompassed by Formula I, include, but are not limited to the compounds in Table I and their pharmaceutically acceptable acid addition salts. In addition, if the compound of the invention is obtained as an acid addition salt, the free base can be obtained by basifying a solution of the acid salt. Conversely, if the product is a free base, an addition salt, particularly a pharmaceutically acceptable addition salt, may be produced by dissolving the free base in a suitable organic solvent and treating the solution with an acid, in accordance with conventional procedures for preparing acid addition salts from base compounds.

Non-toxic pharmaceutical salts include salts of acids such as hydrochloric, phosphoric, hydrobromic, sulfuric, sulfinic, formic, toluenesulfonic, methanesulfonic, nitric, benzoic, citric, tartaric, maleic, hydroiodic, alkanolic such as acetic, $\text{HOOC}-(\text{CH}_2)_n-\text{COOH}$ where n is 0-4, and the like. Those skilled in the art will recognize a wide variety of non-toxic pharmaceutically acceptable addition salts.

The present invention also encompasses the acylated prodrugs of the compounds of Formula I. Those skilled in the art will recognize various synthetic methodologies which may be employed to prepare non-toxic pharmaceutically acceptable addition salts and acylated prodrugs of the compounds encompassed by Formula I.

By "alkyl", "lower alkyl", or $\text{C}_1\text{-C}_6$ alkyl in the present invention is meant straight or branched chain alkyl groups having 1-6 carbon atoms optionally forming a 3 to 6 atoms carbocycle, such as, for example, methyl, ethyl, propyl, isopropyl, cyclopropyl, cyclopropylmethyl, n-butyl, sec-butyl, tert-butyl, cyclobutyl, pentyl, 2-pentyl, isopentyl, neopentyl,


cyclopentyl, hexyl, 2-hexyl, 3-hexyl, 3-methylpentyl, cyclohexyl.

By C₀-C₆ alkylene is meant a direct bond or a C₁-C₆ alkylene group, optionally forming a 3 to 6 atoms carbocycle, such as methylene, ethylidene, propylidene, butylidene, pentylidene, cyclopentylidene, hexylidene, cyclohexylidene.

By "alkoxy", "lower alkoxy", or C₁-C₆ alkoxy in the present invention is meant straight or branched chain alkoxy groups having 1-6 carbon atoms optionally forming a 3 to 6 atoms carbocycle, such as, for example, methoxy, ethoxy, propoxy, isopropoxy, cyclopropylmethoxy, n-butoxy, sec-butoxy, tert-butoxy, pentoxy, 2-pentoxy, isopentoxy, neopentoxy, cyclopentoxy, hexoxy, 2-hexoxy, 3-hexoxy, 3-methylpentoxy, cyclohexoxy.

By "alkanoyl", "lower alkanoyl", or C₁-C₆ alkanoyl as used herein is meant straight or branched chain alkanoyl groups having 1-6 carbon atoms optionally forming a 3 to 6 atom carbocycle, such as, for example, acetyl, propionyl, isopropionyl, cyclopropionyl, butanoyl, pentanoyl, cyclopentanoyl, hexanoyl, cyclhexanoyl. The "ω-position" of the alkanoyl groups herein is the terminal carbon atom.

CONH represents an amide functional group, i.e.,



The term "heterocycle" or "heterocycloalkyl" means a monocyclic or bicyclic hydrocarbon group which in which one or more of the ring carbon atoms has been replaced with a heteroatom, e.g., oxygen, sulfur or nitrogen. Such groups preferably have 4 to 10 carbon atoms and 1 to 4 heteroatoms.

By the term "halogen" in the present invention is meant fluorine, bromine, chlorine, and iodine.

Representative aminoalkyl substituted 9H-pyridino[2,3-b]indole and 9H-pyrimidino[4,5-b]indole derivatives of the present invention, which are encompassed by Formula I, include, but are not limited to the compounds in Examples 1-18 and their
5 pharmaceutically acceptable addition salts.

The interaction of aminoalkyl substituted 9H-pyridino[2,3-b]indole and 9H-pyrimidino[4,5-b]indole derivatives of the invention with CRF₁ and NPY₁ receptors is shown in the examples. This interaction results in the pharmacological
10 activities of these compounds as illustrated in relevant animal models.

The compounds of general Formula I may be administered orally, topically, parenterally, by inhalation or spray or rectally in dosage unit formulations containing conventional
15 non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques. In addition, there is provided a pharmaceutical formulation comprising a
20 compound of general Formula I and a pharmaceutically acceptable carrier. One or more compounds of general Formula I may be present in association with one or more non-toxic pharmaceutically acceptable carriers and/or diluents and/or adjuvants and if desired other active ingredients. The
25 pharmaceutical compositions containing compounds of general Formula I may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsion, hard or soft capsules, or syrups or elixirs.

30 Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and

preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydropropylmethylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example, lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters

derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

Oily suspensions may be formulated by suspending the active ingredients in a vegetable oil, for example arachid oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide palatable oral preparations. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

Pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachid oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol, anhydrides, for example sorbitan monooleate, and condensation

products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavoring agents.

Syrups and elixirs may be formulated with sweetening
5 agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents. The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleaginous suspension. This suspension
10 may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be sterile injectable solution or suspension in a non-toxic parentally acceptable diluent or
15 solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland
20 fixed oil may be employed including synthetic mono-or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

The compounds of general Formula I may also be administered in the form of suppositories for rectal
25 administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials are cocoa butter and
30 polyethylene glycols.

Compounds of general Formula I may be administered parenterally in a sterile medium. The drug, depending on the vehicle and concentration used, can either be suspended or dissolved in the vehicle. Advantageously, adjuvants such as

local anesthetics, preservatives and buffering agents can be dissolved in the vehicle.


Dosage levels of the order of from about 0.1Ümg to about 140Ümg per kilogram of body weight per day are useful in the treatment of the above-indicated conditions (about 0.5Ümg to about 7Üg per patient per day). The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. Dosage unit forms will generally contain between from about 1Ümg to about 500Ümg of an active ingredient.

It will be understood, however, that the specific dose level for any particular patient will depend upon a variety of factors including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, route of administration, and rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

Preparation of Aminoalkyl Substituted 9H-Pyridino[2,3-b]indole and 9H-Pyrimidino[4,5-b]indole Analogues


An illustration of the preparation of compounds of the present invention is given in Scheme I, Scheme II and Scheme III. Those having skill in the art will recognize that the starting materials may be varied and additional steps employed to produce compounds encompassed by the present invention.

Scheme I




wherein Ar, R^1 and R^3 are as defined above for Formula I;
 5 and R^{14} , R^{15} and R^{16} are encompassed by the definition of X as
 defined in Formula I.

Scheme II



wherein Ar , R^1 and R^3 are as defined above for Formula I;
and R^{15} and R^{16} are encompassed by the definition of X as
5 defined in Formula I.

Scheme III



wherein Ar and R¹ are as defined as above for Formula I;
 and R¹⁵, R¹⁶ and R¹⁷ are encompassed by the definition of X as
 5 defined in Formula I.

The disclosures of all articles and references mentioned
 in this application, including patents, are incorporated herein
 by reference.

10 The preparation of the compounds of the present invention
 is illustrated further by the following Examples, which are not
 to be construed as limiting the invention in scope or spirit to
 the specific procedures and compounds described in them.


Commercial reagents were used without further
 purification. THF refers to tetrahydrofuran. LDA refers to
 15 lithium diisopropylamide and DDQ refers to 2,3-dichloro-5,6-

dicyano-1,4-benzoquinone. Room or ambient temperature refers to 20 to 25°C. Concentration implies the use of a rotary evaporator. TLC refers to thin layer chromatography. Mass spectral data were obtained either by CI or APCI methods.

5

Example 1


A. 2-Amino-4,5,6,7-tetrahydro-1-phenyl-1H-indole-3-carbonitrile



- 10 A mixture of 2,4,6-trimethylaniline (500 g) and adipoin (464 g) in toluene (2.5 L) is heated to reflux. A theoretical amount of water is removed azeotropically over the course of 3 hours. The mixture is cooled to ambient temperature, then malononitrile (244 g) and ammonium acetate (57 g) are added.
- 15 The reaction is slowly re-heated back to reflux for about 1 hour with azeotropic removal of water. After cooling, the precipitate that forms overnight is collected by filtration. The dark solid is washed with ethanol and dried to afford 540 g of a white powder: MS 280 (M+H).


20

B. 4-Amino-2-methyl-9-(2,4,6-trimethylphenyl)-5,6,7,8-tetrahydro-9H-pyridino[2,3-b]indole



To the product of Example 1A (535 g) dissolved in
5 dichloroethane (4 L) are added 2-methoxypropene (550 mL) and p-
toluenesulfonic acid monohydrate (3.6 g). The mixture is
refluxed for 1 hour then the solvent is removed by
distillation. The residue is dissolved in THF (3 L) and cooled
to 0°C. To this solution, under an atmosphere of nitrogen gas,
10 is added LDA (2.0M, 1.2 L) at a rate to keep the reaction's
internal temperature below 10°C. After 3 hours the reaction is
neutralized with aqueous HCl. The aqueous layer is extracted
with ethyl acetate and combined with the THF layer. The
combined organic phase is extracted with 3M HCl and the latter
15 is made alkaline (pH=10) with 10N NaOH and ice. The aqueous
solution is extracted with dichloromethane, dried with sodium
sulfate, filtered and concentrated to give a crystalline solid:
MS 320 (M+H).


C. 4-Amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole



The product of Example 1B (600 g) is dissolved in
5 decahydronaphthalene (4 L) and heated to distill off low
boiling impurities that are present. The solution is cooled to
ambient temperature and charged with 10% Palladium on Carbon
(250 g) under a blanket of argon gas. The mixture is heated to
the reflux temperature of 191-193°C for 9 hours to afford
10 aromatized product. The cooled mixture is diluted with
dichloromethane and filtered through a pad of celite. The
dichloromethane in the filtrate is removed under reduced
vacuum. The remaining decahydronaphthalene solution is treated
by bubbling in a stream of hydrochloric acid gas with ice-
15 cooling for about 5 minutes. The solids are filtered, washed
with diethyl ether and dried to yield 580 g of product as the
HCl salt. The salt is recrystallized from an ethyl acetate and
ethanol mixture to give a white product: MS(free base) 316
(M+H).

20


D. 4-(N-Cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole



A solution of dichloroethane (250 mL) containing the
5 product of Example 1C (50 g as HCl salt) and
cyclopropanecarbonyl chloride (14 mL) at reflux is treated with
dropwise addition of N,N-diisopropylethylamine (54 mL). After
heating for 0.5 hour the reaction is cooled to ambient
temperature and poured into aqueous potassium carbonate
10 solution. The product is extracted with dichloromethane, dried
over sodium sulfate, filtered and concentrated to give a white
solid. This solid is re-dissolved in THF (600 mL) and mixed
with borane-methyl sulfide complex (10M, 43 mL). The mixture
is heated to reflux for 8 hours and quenched at room
15 temperature with a large excess of methanol (about 200 mL).
Re-heat mixture to reflux for 1 hour, then concentrate under
reduced pressure. More methanol (another 200 mL) is added to
the gummy residue and the solution is re-concentrated to yield
a white solid: MS 370 (M+H).


20

E. 4-(N-(2-Chloroethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole



A solution containing the product from Example 1D (52 g)
5 and chloroacetyl chloride (34 mL) in dichloroethane (500 mL)
is refluxed for 4 hours. The solvent and excess reagent are
removed under reduced pressure. Aqueous potassium carbonate
is added to the remaining oily residue and extracted with
dichloromethane. The extract is dried with sodium sulfate,
10 filtered and concentrated. The latter chloroacetyl compound
(63 g) is dissolved in THF (250 mL). Add borane-methyl
sulfide complex (10M, 14 mL) and stir at ambient temperature
for 15 minutes then for 1 hour at reflux temperature. The
solution is cooled back to room temperature, quenched with a
15 large excess of methanol (100 mL) and re-heated to reflux for
1 hour. The solution is concentrated to give a viscous oil
that crystallizes on standing: MS 432 (M+H).


F. 4-(N-(2-Pyrrolidinoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole (Compound 1)



5 A steel bomb containing the product from Example 1E (61 g), pyrrolidine (60 mL) and N-methylpyrrolidinone (250 mL) is sealed and heated to 120°C for 5 hours. The mixture is poured into water and extracted with ethyl acetate. The organic layer is washed with water, dried over sodium sulfate,
10 filtered and concentrated. The product is purified by flash chromatography. First 100% ethyl acetate is used to elute impurities followed by 10% methanol in dichloromethane to elute the desired product. Obtain 60 g of compound that crystallizes on standing. The sulfate salt of the amine is
15 formed by adding concentrated sulfuric acid (2.44 mL) to amine (21.4 g) dissolved in ethanol (50 mL). Isopropanol (200 mL) is added to the ethanol solution at reflux. Crystals of sulfate that form on cooling are collected by filtration and washed with cold isopropanol: MS(free base) 467 (M+H).


20

G. 4-Propylamino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole



An excess of sodium borohydride (about 1 g) is added in
5 small portions to a propionic acid (10 mL) solution of the
product from Example 1C (1.4 g). After gas evolution ceases
the reaction is heated to 100°C for 0.5 hour. Propionic acid
is removed from the mixture under reduced pressure and crude
product is extracted from aqueous potassium carbonate with
10 dichloromethane. The dichloromethane extract is dried over
sodium sulfate, filtered and concentrated: MS 358 (M+H).

H. 4-Cyclopropylamino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole



To a solution of the product from Example 1C (10 g) and 2-methoxypropene (50 mL) in dichloroethane (100 mL) is added acetic acid (1.6 mL) and sodium triacetoxyborohydride (25 g). The mixture is stirred at room temperature for 24 hours then
5 concentrated. Dissolve residue in ethyl acetate and wash with water, followed by 1N sodium hydroxide and brine. Dry over sodium sulfate, filter and concentrate: MS 358 (M+H).

Example 2

10. The following compounds are prepared essentially according to the procedures set forth above in Example 1.

a) 4-(N-(2-N',N'-Dimethylaminoethyl)-N-propyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 429 (M+H). (Compound 2)

15

b) 4-(N-(2-N',N'-Dimethylaminoethyl)-N-2-methoxyethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 445 (M+H). (Compound 3)

20

c) 4-(N-(2-N',N'-Dimethylaminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 441 (M+H). (Compound 4)

d) 4-(N-(2-N'-Ethylaminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 441 (M+H). (Compound 5)

e) 4-(N-(2-(N'-Ethyl-N'-methyl)aminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 455 (M+H). (Compound 6)

f) 4-(N-2-(2-(S)-Methoxymethylpyrrolidino)ethyl-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 511 (M+H). (Compound 7)

g) 4-(N-(2-N',N'-Dimethylaminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(4-methoxy-2-methylphenyl)-9H-pyridino[2,3-b]indole: MS 443 (M+H). (Compound 8)

5

h) 4-(N-(2-N',N'-Diethylaminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 469 (M+H). (Compound 9)

10

i) 4-(N-(2-N'-Methylaminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 427 (M+H). (Compound 10)

15

j) 4-(N-(2-Piperidinoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 481 (M+H). (Compound 11)

20

k) 4-(N-(2-Pyrrolidinoethyl)-N-propyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 455 (M+H). (Compound 12)

25

l) 4-(N-(2-Piperidinoethyl)-N-propyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 469 (M+H). (Compound 13)

30

m) 4-(N-(2-Morpholinoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 483 (M+H). (Compound 14)

n) 4-(N-(2-N'-Ethyl-N-methylaminoethyl)-N-propyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 443 (M+H). (Compound 15)

o) 4-(N-(2-(1-Imidazolyl)ethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 464 (M+H). (Compound 16)

5 p) 4-(N-2-(N'-Methylpiperazinyl)ethyl-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 496 (M+H). (Compound 17)

q) 4-(N-2-(N'-Methylhomopiperazinyl)ethyl-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 510 (M+H). (Compound 18)

r) 4-(N-(2-N'-Isopropylaminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 454 (M+H). (Compound 19)

s) 4-(N-(2-(2-(1-Methyl-2-pyrrolidino)ethyl)aminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 524 (M+H). (Compound 20)

20 t) 4-(N-(2-Piperazinylethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 482 (M+H). (Compound 21)

u) 4-(N-(2-Pyrrolidinoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4-dimethylphenyl)-9H-pyridino[2,3-b]indole: MS 453 (M+H). (Compound 22)

v) 4-(N-(3-Pyrrolidinopropyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 481 (M+H). (Compound 23)

w) 4-(N-(2-Methyl-2-pyrrolidinoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 481 (M+H). (Compound 24)

x) 4-(N-(4-Pyrrolidinobutyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 495 (M+H). (Compound 25)

5

y) 4-(N-(2-(4-Piperidinopiperidinyl)ethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 564 (M+H). (Compound 26)

10

z) 4-(N-(2-(2-Phenethylamino)ethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 517 (M+H). (Compound 27)

15

aa) 4-(N-(2-N'-Methylaminoethyl)-N-methyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 387 (M+H). (Compound 28)

20

bb) 4-(N-(2-Pyrrolidinoethyl)-N-2-methoxyethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 471 (M+H). (Compound 29)

25

cc) 4-(N-(2-Pyrrolidinoethyl)-N-ethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 441 (M+H). (Compound 30)

dd) 4-(N-(2-Pyrrolidinoethyl)-N-butyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 469 (M+H). (Compound 31)

30

ee) 4-(N-(2-Pyrrolidinoethyl)-N-2-methylpropyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 469 (M+H). (Compound 32)

ff) 4-(N-(2-N',N'-Dimethylaminoethyl)-N-2-methylpropyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 443 (M+H). (Compound 33)

5 gg) 4-(N-(2-(N'-Ethyl-N-methyl)aminoethyl)-N-2-methylpropyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 457 (M+H). (Compound 34)

hh) 4-(N-(2-Pyrrolidinoethyl)-N-isopropyl)amino-2-methyl-10 9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 455 (M+H). (Compound 35)

ii) 4-(N-(2-N',N'-Dimethylaminoethyl)-N-isopropyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: 15 MS 429 (M+H). (Compound 36)

jj) 4-(N-(2-Guanidinoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 455 (M+H). (Compound 37)

20 kk) 4-(N-(2-(2-(4-Methoxy)phenethylamino)ethyl)-N-ethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 521 (M+H). (Compound 38)


ll) 4-(N-(2-N'-Cyclopentylaminoethyl)-N-ethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 25 455 (M+H). (Compound 39)

mm) 4-(N-(2-N',N'-Dimethylaminoethyl)-N-ethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 30 415 (M+H). (Compound 40)

nn) 4-(N-(2-N',N'-Dimethylaminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,6-dimethylphenyl)-9H-pyridino[2,3-b]indole: MS 427 (M+H). (Compound 41)

Example 3

4-(N-(2-aminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-
(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole (Compound
5 42)



A solution of the product from Example 1E (900 mg) and sodium azide (410 mg) in N-methylpyrrolidinone (10 mL) is heated to 120°C for 2 hours. The mixture is poured into water and extracted with ethyl acetate. The organic layer is washed with water, dried over sodium sulfate, filtered and concentrated. An ethanol (10 mL) solution of the crude product and 10% palladium on carbon (about 300 mg) is hydrogenated for 8 hours at approximately 1 atmosphere pressure. The suspension is filtered over celite and the concentrated product is purified by flash chromatography, then converted to the hydrochloride salt: MS (free base) 413 (M+H). (Compound 42)

The following compounds are prepared essentially according to the procedure set forth in Example 3.

a) 4-(N-(3-Aminopropyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 427 (M+H). (Compound 43)

b) 4-(N-(2-Amino-2-methylethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 427 (M+H). (Compound 44)

5

c) 4-(N-(2-Aminoethyl)-N-propyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 401 (M+H). (Compound 45)

10

d) 4-(N-(2-Aminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4-dimethylphenyl)-9H-pyridino[2,3-b]indole: MS 399 (M+H). (Compound 46)

15

e) 4-(N-(2-Aminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,6-dimethylphenyl)-9H-pyridino[2,3-b]indole: MS 399 (M+H). (Compound 47)


20

f) 4-(N-(2-Aminoethyl)-N-ethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 387 (M+H). (Compound 48)

g) 4-(N-(4-Aminobutyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 441 (M+H). (Compound 49)

Example 4

4-(N-(2-(4-Triazolyl)ethyl)-N-cyclopropylmethyl)amino -2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole
(Compound 50)



5


A solution containing the product from Example 3 (380 mg) in toluene (8 mL) with N,N'-dimethylformamide azine (195 mg) and p-toluenesulfonic acid monohydrate (6 mg) is heated at reflux. A flowing stream of nitrogen gas is used to displace evolving dimethylamine. After 12 hours heating is discontinued and the solution is diluted with ethyl acetate and washed with water, brine and dried over sodium sulfate, filtered and concentrated to a tan colored solid. The product is purified by preparative TLC using 7% methanol and 0.1% ammonium hydroxide in dichloromethane as eluent to afford 340 mg of product: MS 465 (M+H).

10

15

Example 5

4-(N-(2-Carboxamidoethyl)-N-cyclopropylmethyl)amino-2-
methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole
5 (Compound 51)




Dissolve the compound from Example 3 (1.0 g) and sodium cyanide (540 mg) in N-methylpyrrolidinone (20 mL) and heat to 95°C for 2 hours. The mixture is poured into water and
10 extracted with ethyl acetate. The organic layer is washed with water, dried over sodium sulfate, filtered and concentrated. Purify by flash chromatography using 20% ethyl acetate in hexanes to yield a white solid. Dissolve nitrile (130 mg) and solid sodium hydroxide (800 mg) in tert-butanol (5 mL). Reflux
15 mixture for about 1 hour. Next, neutralize with 6N hydrochloric acid, concentrate and purify residue by preparative TLC using 10% methanol in dichloromethane as eluent: MS 441(M+H).

20

Example 6

4-(N-(2-N'-acetyl-N'-methylaminoethyl)-N-
cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-
pyridino[2,3-b]indole (Compound 53)



To a solution of the compound from Example 2i (170 mg) in dichloromethane (5 mL) is added acetic anhydride (0.1 mL) and diisopropylethylamine (0.2 mL). The solution is stirred for 0.5 hour then concentrated. The residue is dissolved in ethyl acetate, washed with sodium carbonate and water, dried over sodium sulfate, filtered and concentrated. The product is purified by preparative TLC using 70% ethyl acetate in hexanes as eluent: MS 469 (M+H).

10

Example 7

The following compounds are prepared essentially according to the procedures set forth above in Example 6.

a) 4-(N-(2-(N'-Methanesulfonyl-N-methylamino)ethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 505 (M+H). (Compound 54)

b) 4-(N-(2-(N'-Methyl-N'-trifluoromethanesulfonyl)aminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 559 (M+H). (Compound 55)

c) 4-(N-(2-(4-Aminophenylsulfonamido)ethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 568 (M+H). (Compound 56)


5 d) 4-(N-(2-(2-Thienylsulfonamido)ethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 559 (M+H). (Compound 57)

10 e) 4-(N-(2-(2-(2-Thienyl)-1-oxoethyl)aminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 537 (M+H). (Compound 58)

15 f) 4-(N-(2-(2-Phenyl-1-oxoethyl)aminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 531 (M+H). (Compound 59)

Example 8

20 4-(N-(2-N',N'-dimethylamino-1-oxoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole (Compound 60)



Refrigerated dimethylamine (about 4 mL) is poured into a solution of the chloroacetyl compound from Example 1E (1.9 g)

and acetonitrile (20 mL). The reaction vessel is sealed and the mixture is stirred at ambient temperature for 2 hours then concentrated. Water is added to the residue and the aqueous is extracted with ethyl acetate. The extract is dried over sodium sulfate, filtered and concentrated. The product is purified on preparative TLC with 100% ethyl acetate as the eluting solvent: MS 443 (M+H).

Example 9

The following compounds are prepared essentially according to the procedure set forth in Example 8.

a) 4-(N-(2-N',N'-Dimethylamino-1-oxoethyl)-N-propyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 443 (M+H). (Compound 61)

b) 4-(N-(2-N',N'-Dimethylamino-1-oxoethyl)-N-2-methoxyethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 459 (M+H). (Compound 62)

c) 4-(N-(2-Pyrrolidino-1-oxoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 481 (M+H). (Compound 63)

d) 4-(N-(2-N'-Ethylamino-1-oxoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 455 (M+H). (Compound 64)


e) 4-(N-(2-(N'-Ethyl-Nⁱ-methyl)amino-1-oxoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 469 (M+H). (Compound 65)

f) 4-(N-(2-(2-(S)-Methoxymethylpyrrolidino)-1-oxoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 525 (M+H). (Compound 66)

5 g) 4-(N-(2-(1-Imidazolyl)-1-oxoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 478 (M+H). (Compound 67)


Example 10

1.0 A. 4-(N-(2-Chloro-1-oxoethyl)-N-propyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole



A solution of the compound from Example 1C (5.7 g),
15 chloroacetyl chloride (3 mL) and diisopropylethylamine (3 mL)
in dichloroethane (90 mL) is refluxed for 0.5 hour. After
concentrating the mixture, aqueous potassium carbonate is
added and product is extracted with dichloromethane. The
extract is dried over sodium sulfate, filtered and
20 concentrated. The chloride (1.0 g) is dissolved in
acetonitrile (20 mL) and mixed with pyrrolidine (2 mL). The
solution is stirred for 2 hours at ambient temperature then
concentrated. Water is added to the residue and the aqueous
is extracted with ethyl acetate. The extract is dried over
25 sodium sulfate, filtered and concentrated: MS 392 (M+H).

B. 4-(N-(2-pyrrolidinoethyl)-N-1-oxopropyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole
(Compound 68)



5 To a solution of the compound from Example 10A (1.0 g) in acetonitrile (25 mL) is added pyrrolidine (1.0 mL). The solution is stirred at room temperature for 2 hours then concentrated. Water is added to the residue and the aqueous is extracted with ethyl acetate. The extract is dried over sodium
10 sulfate, filtered and concentrated. The isolated aminoamide (1.1 g) is dissolved in THF (25 mL), treated with borane-methyl sulfide complex (10M, 1.0 mL) and refluxed for 8 hours. The solution is cooled and quenched with an excess of methanol (25 mL). Next, N,N'-dimethylethylenediamine (1.1 mL) is added and
15 the mixture is re-heated to reflux for 4 hours. The solution is concentrated, diluted with ethyl acetate, washed with water, dried over sodium sulfate, filtered and concentrated. To a solution containing the latter product (330 mg), diisopropylethylamine (0.2 mL) and dichloroethane (10 mL) is
20 added propionyl chloride (0.1 mL). The mixture is refluxed for 0.5 hour, then poured into aqueous potassium carbonate and extracted with ethyl acetate. The extract is dried over sodium sulfate, filtered and concentrated. The product is purified on

preparative TLC with 10% methanol in dichloromethane as the eluting solvent: 469 (M+H).

Example 11

5 The following compounds are prepared essentially according to the procedure set forth in Example 10.

a) 4-(N-(2-Pyrrolidinoethyl)-N-cyclopropyloxomethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 481 (M+H). (Compound 69)


10

b) 4-(N-(2-Dimethylaminoethyl)-N-methoxymethyloxomethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 459 (M+H). (Compound 70)

15

Example 12

4-(N-(2-pyridylmethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole (Compound 71)



20

A solution of the compound from Example 1D (110 mg), picolinyl chloride (98 mg) and N,N-diisopropylethylamine in DMF (5 mL) is stirred at 60°C for 12 hours. The mixture is

poured into water and extracted with ethyl acetate. The extract is washed with water, dried over sodium sulfate, filtered and concentrated. The product is purified by preparative TLC using 30% ethyl acetate in hexanes as eluent:
5 MS 461 (M+H).

Example 13

The following compounds are prepared essentially according to the procedure of Example 12.

10 a) 4-(N-(2-N',N'-Dimethylamino-2-oxoethyl)-N-2-methoxyethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 459 (M+H). (Compound 72)

b) 4-(N-(2-N',N'-Dimethylamino-2-oxoethyl)-N-propyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 443 (M+H). (Compound 73)


c) 4-(N-(2-N',N'-Dimethylamino-2-oxoethyl)-N-2-methoxy-1-oxoethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 473 (M+H). (Compound 74)

d) 4-(N-(2-N',N'-Dimethylamino-1-oxoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 455 (M+H). (Compound 75)

25

Example 14


4-(4-Triazolyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole (Compound 76)



A solution containing the compound from Example 1C (560 mg) in toluene (10 mL) with N,N-dimethylformamide azine (400 mg) and p-toluenesulfonic acid monohydrate (50 mg) is heated at reflux. A flowing stream of nitrogen gas is used to displace evolving dimethylamine. After 24 hours heating is discontinued and the solution is diluted with ethyl acetate and washed with water, brine and dried over sodium sulfate, filtered and concentrated to a tan colored solid. The product is purified by preparative TLC using 100% ethyl acetate as eluent to afford 340 mg of product: MS 368 (M+H).


Example 15

15 A. 4-Chloro-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole



Dissolve tert-butyl nitrite (0.65 g) in acetonitrile (10 mL) and add copper(II) chloride (0.68 g). Then the compound from Example 1C (1.33 g) is added portionwise to the greenish-brown solution and the mixture is stirred for 12 hours. The
5 acetonitrile is removed by evaporation and the residue is partitioned between water and dichloromethane. The aqueous layer is extracted with more dichloromethane and the combined extract is washed with water, dried over sodium sulfate, filtered and concentrated. The product is filtered through a
10 plug of silica gel using 20% ethyl acetate in hexanes as eluent to afford a tan colored solid: MS 335 (M+H).

B. 4-Piperazinyl-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole (Compound 77)



Combine the compound from Example 15A (200 mg) and
5 piperazine (0.58 g) in N-methylpyrrolidinone (2 mL) and heat
the solution to 120 °C for 12 hours. Pour mixture into water
and extract with ethyl acetate. Wash extract with aqueous
ammonium chloride then water. Dry extract over sodium sulfate,
filter and concentrate. Purify by preparative TLC using 10%
10 methanol in dichloromethane as eluent: MS 385 (M+H).

Example 16

The following compounds are prepared essentially
15 according to the procedures set forth above in Example 15.

a). 4-(4-Methylpiperazinyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 399 (M+H).
(Compound 78)

20

b) 4-Homopiperazinyl-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 399 (M+H). (Compound 79)

c) 4-(N-(2-(1-Methyl-2-pyrrolidino)ethyl)-N-cyclopropylcarbonyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 495 (M+H). (Compound 80)

5 d) 4-(N-2-(S)-Pyrrolidinomethylpyrrolidinyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 453 (M+H). (Compound 81)

10 e) 4-(4-Aminomethylpiperidino)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 413 (M+H). (Compound 82)

15 f) 4-(4-Piperidinopiperidinyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 467 (M+H). (Compound 83)

20 g) 4-(4-Carboxamidopiperidinyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 427 (M+H). (Compound 84)

h) 4-(3-Aminopyrrolidinyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 385 (M+H). (Compound 85)

25 i) 4-(N-(2-(1-Methyl-2-pyrrolidino)ethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 481 (M+H). (Compound 86)

30 j) 4-(N-2-(S)-Cyclopentylaminomethylpyrrolidinyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 467 (M+H). (Compound 87)

k) 4-(N-2-(R)-Cyclopentylaminomethylpyrrolidinyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 467 (M+H). (Compound 88)

5 l) 4-(N-3-Cyclopentylaminopyrrolidinyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 467 (M+H). (Compound 89)

10 m) 4-(N-3-(2-(3-Methoxy-4-ethoxy)phenethyl)aminopyrrolidinyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 563 (M+H). (Compound 90)

15 n) 4-(N-3-(1-oxo-2-(3-Methoxy-4-ethoxy)phenethyl)aminopyrrolidinyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 577 (M+H). (Compound 91)


20 o) 4-(N-2-(2-(R)-(4-Methoxy)phenethyl)aminomethylpyrrolidinyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 533 (M+H). (Compound 92)

p) 4-(N-2-(2-(S)-(4-Methoxy)phenethyl)aminomethylpyrrolidinyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 533 (M+H). (Compound 93)

25 q) 4-(N-4-(2-(3-Methoxy-4-ethoxy)phenethyl)aminomethylpiperazinyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole: MS 563 (M+H). (Compound 94)


Example 17

30 A. 2-Amino-1-phenyl-1H-indole-3-carbonitrile



Dissolve the compound from Example 1A (20 g) in 1,4-dioxane (300 mL) and add DDQ (34 g) portionwise to the solution. The reaction is stirred for 1 hour then filtered
5 through celite to remove insoluble side products. The filtrate is concentrated and allowed to solidify. The product is collected by filtration and washed with ethanol to yield 16 g of a tan colored powder: MS 276 (M+H).


10 B. 4-Hydroxy-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole



15 A mixture of the compound from Example 12A (30 g), acetic anhydride (15 mL) and acetic acid (30 mL) is refluxed for 1 hour then, concentrated to a solid. Phosphoric acid (40 mL, 85%) is added to the amide. The mixture is then refluxed for 0.5 hour and cooled to ambient temperature. The solution
20 is poured onto ice and the precipitate that forms is collected

by filtration. The solids are washed with water and some ethanol: MS 318 (M+H).


5 C. 4-Chloro-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole



The compound from Example 12B (2.2 g) is refluxed in phosphoryl chloride (30 mL) for 3 hours. The excess phosphoryl chloride is removed under reduced pressure and the residue is partitioned between aqueous potassium carbonate and dichloromethane. The aqueous is extracted with more dichloromethane. The combined extracts are dried over sodium sulfate, filtered and concentrated to give a tan colored solid: MS 336 (M+H).


15

D. 4-Cyclopropylamino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole



A mixture of the compound from Example 12C (750 mg) and cyclopropylamine (1.6 mL) in N-methylpyrrolidinone (2 mL) is heated to 65°C in a sealed tube for 24 hours. Dilute mixture with ethyl acetate and wash with water, brine, dry over sodium sulfate, filter and concentrate to give a tan colored solid. Purify by radial chromatography using 40% ethyl acetate in hexanes as eluent to give 730 mg of product: MS 357 (M+H).

E. 4-(N-(2-N',N'-Dimethylaminoethyl)-N-cyclopropyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole (Compound 95)



To a solution of compound from Example 12D (200 mg) in dimethylformamide (5 mL) at 0°C, under a blanket of nitrogen, is added sodium hydride (60%, 100 mg). After stirring the solution for 15 minutes, 2-dimethylaminoethyl chloride hydrochloride (170 mg) is added. The mixture is then heated to 60°C for 2 hours, then quenched with ice and water. Diluted with ethyl acetate and washed with water, brine, dried over sodium sulfate, filtered and concentrated. Purified by radial chromatography using 7% methanol and 0.5% ammonium hydroxide in dichloromethane as eluent to obtain 190 mg of product: MS 428 (M+H).

Example 18

The following compounds are prepared essentially according to the procedure of Example 17.

a) 4-(N-(2-N',N'-Dimethylaminoethyl)-N-methyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 466 (M+H). (Compound 96)

b) 4-(N-(2-N',N'-Dimethylaminoethyl)-N-ethyl)amino-2-methyl-9-(4-bromo-2,6-dimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 480 (M+H). (Compound 97)

c) 4-(N-(2-N',N'-Dimethylaminoethyl)-N-ethyl)amino-2-methyl-9-(4-chloro-2-methylphenyl)-9H-pyrimidino[4,5-b]indole: MS 422 (M+H). (Compound 98)

d) 4-(N-(2-Morpholinoethyl)-N-propyl)amino-2-methyl-9-(2-chloro-4-methylphenyl)-9H-pyrimidino[4,5-b]indole: MS 478 (M+H). (Compound 99)

e) 4-(N-(2-N',N'-Dimethylaminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 442 (M+H). (Compound 100)

f) 4-Piperazinyl-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 386 (M+H). (Compound 101)

g) 4-(N-(2-Pyrrolidinoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 468 (M+H). (Compound 102)

h) 4-(3-Aminopyrrolidinyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 386 (M+H). (Compound 103)

i) 4-(2-(S)-Aminomethylpyrrolidino)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 400 (M+H).
(Compound 104)

5

j) 4-(3-Amino-2-triazolyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 384 (M+H).
(Compound 105)

10

k) 4-(3-Amino-1-triazolyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 384 (M+H).
(Compound 106)

15

l) 4-(3-Amino-4-triazolyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 384 (M+H).
(Compound 107)

20

m) 4-(N-(2-Pyrrolidinoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(4-bromo-2,6-dimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 532 (M+H). (Compound 108)

25

n) 4-(N-(2-Pyrrolidinoethyl)-N-propyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 456 (M+H). (Compound 109)

o) 4-(N-(2-Pyrrolidinoethyl)-N-butyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 470 (M+H). (Compound 110)

30

p) 4-(N-(2-Pyrrolidinoethyl)-N-cyclopropyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 454 (M+H). (Compound 111)

q) 4-(N-(2-N',N'-Dimethylaminoethyl)-N-propyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole:
MS 430 (M+H). (Compound 112)

5 r) 4-(N-(2-N',N'-Dimethylaminoethyl)-N-butyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole:
MS 444 (M+H). (Compound 113)

s) 4-(N-(2-Piperidinoethyl)-N-propyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 470
10 (M+H). (Compound 114)

t) 4-(N-(2-Piperidinoethyl)-N-butyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 484
15 (M+H). (Compound 115)

u) 4-(N-(2-N',N'-Dimethylaminoethyl)-N-cyclobutyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole:
MS 442 (M+H). (Compound 116)

20

v) 4-(N-(2-N',N'-Dimethylaminoethyl)-N-cyclopentyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole:
MS 456 (M+H). (Compound 117)

25 w) 4-(N-(2-N',N'-Dimethylaminoethyl)-N-isopropyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole:
MS 430 (M+H). (Compound 118)

x) 4-(N-(2-N'-Methylaminoethyl)-N-methyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 388
30 (M+H). (Compound 119)

y) 4-(4-(N-Methylamino)piperidino)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 414 (M+H).
(Compound 120)

5 z) 4-(4-(2-Aminoethyl)piperazinyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 429 (M+H).
(Compound 121)

10 aa) 4-(N-(2-N'-Ethylaminoethyl)-N-ethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 416
(M+H). (Compound 122)

15 bb) (\pm)-4-(N-(2-N',N'-Dimethylaminoethyl)-N-1-methylpropyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 444 (M+H). (Compound 123)

20 cc) 4-(N-(2-N',N'-Diethylaminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 470 (M+H). (Compound 124)

dd) 4-(N-(2-N',N'-Diethylaminoethyl)-N-cyclopropyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 456 (M+H). (Compound 125)

25 ee) 4-(N-(2-N',N'-Dimethylaminoethyl)-N-2-methoxyethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 446 (M+H). (Compound 126)

30 ff) 4-(N-(2-N',N'-Dimethylaminoethyl)-N-2-ethoxyethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 460 (M+H). (Compound 127)

gg) 4-(N-(2-N'-Ethylaminoethyl)-N-ethyl)amino-2-methyl-9-(2-methyl-4-chlorophenyl)-9H-pyrimidino[4,5-b]indole: MS 423 (M+H). (Compound 128)

5 hh) 4-(N-(Cyclopentylaminopiperidin-4-yl)-N-ethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 468 (M+H). (Compound 129)

ii) 4-(N-(2-N',N'-Dimethylaminoethyl)-N-isobutyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 444 (M+H). (Compound 130)

jj) 4-(N-(2-(3-Methoxy-4-ethoxy)phenethyl)aminopiperidin-4-yl)-N-ethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 578 (M+H). (Compound 131)

kk) 4-(N-(S)-(1-Oxo-2-(4-methoxy)phenethyl)amino pyrrolidin-2-yl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 54 (M+H). (Compound 132)

20 ll) 4-(N-(S)-(2-(4-Methoxy)phenethyl)amino pyrrolidin-2-yl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole: MS 578 (M+H). (Compound 133)

Example 19

25 The pharmaceutical utility of compounds of this invention is indicated by the following assays for human CRF1 and NPY1 receptor activity.

Assay for CRF Receptor Binding Activity

30 CRF receptor binding is performed using a modified version of the assay described by Grigoriadis and De Souza (Methods in Neurosciences, Vol. 5, 1991). Membrane pellets containing CRF receptors are re-suspended in 50mM Tris buffer pH 7.7 containing 10 mM MgCl₂ and 2 mM EDTA and centrifuged for 10 minutes at 48000g. Membranes are washed again and brought to a

final concentration of 1500mg/ml in binding buffer (Tris buffer above with 0.1 % BSA, 15 mM bacitracin and 0.01 mg/mL aprotinin). For the binding assay, 100 mL of the membrane preparation is added to 96 well microtube plates containing
5 100 mL of ^{125}I -CRF (SA 2200 Ci/mmol, final concentration of 100 pM) and 50 mL of drug. Binding is carried out at room temperature for 2 hours. Plates are then harvested on a Brandel 96 well cell harvester and filters are counted for gamma emissions on a Wallac 1205 Betaplate liquid
10 scintillation counter. Non specific binding is defined by 1 mM cold CRF. IC_{50} values are calculated with the non-linear curve fitting program RS/1 (BBN Software Products Corp., Cambridge, MA). The binding affinity for the compounds of Formula I expressed as IC_{50} value, generally ranges from about
15 0.5 nanomolar to about 10 micromolar.

Alternatively, the binding activity of the compounds of formula I to the human CRF_1 receptor can be measured as follows:

20 Assay for Human CRF Receptor Binding Activity in IMR32 cells

^{125}I]Sauvagine Binding to CRF_1 Receptors Endogenously Expressed in IMR-32 Cells: IMR-32 human neuroblastoma cells are grown to 80% confluence in EMEM containing Earle's Balanced Salts and 2 mM l-glutamine with 10% FBS, 25 mM HEPES, 1 mM
25 Sodium Pyruvate, and nonessential amino acids. At this time, flasks of cells are treated with 2.5 μM 5-bromo-2'-deoxyuridine (Br-dU) for 10 days. Media is changed every 3-4 days across the 10 day period. Cells are harvested using No-Zyme (JRH Biosciences) and rinsed with PBS. For membrane preparation,
30 cells are homogenized in wash buffer (50 mM Tris HCl, 10 mM MgCl_2 , 2 mM EGTA, pH 7.4) and centrifuged at 48,000 x g for 10 minutes at 4°C. Pellets are re-suspended, homogenized and centrifuged two additional times. The receptor binding assay

is performed using assay buffer (50 mM Tris HCl, 10 mM MgCl₂, 2 mM EGTA, pH 7.4, 0.1% BSA, 0.1 mM bacitracin (22.0mg/100 mL)), 150 µg protein/tube, and [¹²⁵I]Sauvagine (NEN; 100 pM for competition analysis and 10 pM-1 nM for saturation analysis) to
5 yield a final volume of 200 µL. Nonspecific binding is defined using 2 µM r/h CRF or 9-41 alpha-helical CRF. Cells are incubated for 2 hours at room temperature. The assay is terminated by rapid vacuum filtration (Tomtec: Deepwell 3) through GFC filters presoaked in 1% PEI using ice-cold 50 mM
10 Tris HCl and dry thoroughly by air. Specific Binding: 70-80%; K_d (nM): 0.30 nM; B_{max} (fmole/mg protein): 40-50. IC₅₀ values are calculated with the non-linear curve fitting program RS/1 (BBN Software Products Corp., Cambridge, MA).

The binding affinities for the compounds of Formula I
15 towards the CRF₁ receptor are expressed as IC₅₀ values and are less than 10 micromolar.

Assay for Human NPY₁ Receptor Binding Activity

Compounds are assayed for activity using the following
20 method: Baculovirus-infected Sf9 cells expressing recombinant human NPY Y₁ receptors are harvested at 42-48 hours at which time batches of 500 mL of cell suspension are pelleted by centrifugation. Each pellet is re-suspended in 30 mL of lysis buffer (10 mM HEPES, 250 mM sucrose, 0.5 µg/mL leupeptin, 2
25 µg/mL Aprotinin, 200 µM PMSF and 2.5 mM EDTA, pH 7.4) and gently homogenized by 50 strokes using a dounce homogenizer. The homogenate is centrifuged at 4°C for 10 minutes at 536 x g to pellet the nuclei. The supernatant is collected into a fresh tube and centrifuged twice in the same buffer at 48,000 x
30 g for 40 minutes. The final pellet is subsequently re-suspended in 10 mL of PBS containing 5 mM EDTA by dounce homogenization and stored in aliquots at -80°C.

Purified membranes are washed by PBS and re-suspended by gentle pipetting in binding buffer (50 mM Tris(HCl), 5 mM KCl, 120 mM NaCl, 2 mM CaCl₂, 1 mM MgCl₂, 0.1% bovine serum albumin (BSA), pH 7.4). Membranes (5μg) are added to siliconized (Sigmacote, Sigma) polypropylene tubes in addition to 0.050 nM [¹²⁵I]NPY (porcine) for competition analysis or 0.010-0.500 nM [¹²⁵I]NPY (porcine) for saturation analysis. For evaluation of guanine nucleotide effects on receptor affinity, GTP is added at a final concentration of 100 μM. Cold displacers are added at concentrations ranging from 10⁻¹² M to 10⁻⁶ M to yield a final volume of 0.250 mL. Nonspecific binding is determined in the presence of 1 μM NPY (human) and accounts for less than 10% of total binding. Following a 2 hour incubation at room temperature, the reaction is terminated by rapid vacuum filtration. Samples are filtered over presoaked GF/C Whatman filters (1.0% polyethyleneimine for 2 hours) and rinsed 2 times with 5 mL cold binding buffer lacking BSA. Remaining bound radioactivity is measured by gamma counting. To estimate the B_{max}, K_d and K_i, the results of binding experiments are analyzed using SigmaPlot software (Jandel).


The binding affinities for the compounds of Formula I towards the NPY₁ receptor are expressed as IC₅₀ values and are less than 10 micromolar.

The invention and the manner and process of making and using it, are now described in such full, clear, concise and exact terms as to enable any person skilled in the art to which it pertains, to make and use the same. It is to be understood that the foregoing describes preferred embodiments of the present invention and that modifications may be made therein without departing from the spirit or scope of the present invention as set forth in the claims. To particularly point

out and distinctly claim the subject matter regarded as invention, the following claims conclude this specification.

WHAT IS CLAIMED IS:

1. A compound of the formula:



or the pharmaceutically acceptable salts thereof wherein


5 wherein

Ar is phenyl, 1- or 2-naphthyl, 2-, 3-, or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, optionally mono-, di-, or tri-substituted with halogen, trifluoromethyl, hydroxy, amino, lower alkylamino, lower dialkylamino, carboxamido, N-lower alkyl carboxamido, N,N-lower dialkyl carboxamido, C₁-C₆ alkyl, C₁-C₆ alkoxy, with the proviso that at least one of the positions ortho or para to the point of attachment of Ar to the tricyclic ring system is substituted;

10 R¹ is hydrogen, halogen, trifluoromethyl, C₁-C₆ alkyl, or (C₁-C₆ alkyl)-G¹-R² where G¹ is oxygen or sulfur and R² is hydrogen or C₁-C₆ alkyl;

W is N or C-R³ where R³ is hydrogen or C₁-C₆ alkyl; and

X is



(i)

wherein

20 V¹ and V² are CH₂, CO, CS, SO₂ or CH(C₁-C₆ alkyl), with the proviso that both V¹ and V² cannot both be CO, CS or SO₂;

Y¹ and Y² independently represent a bond or C₁-C₆ alkylene;


A¹ is NR⁴R⁵ wherein R⁴ and R⁵ are independently
hydrogen, a lower alkyl group which optionally forms

a heterocycloalkyl group with Y¹;

C₁-C₆ arylalkyl or C₁-C₆ heteroarylalkyl, where aryl
is phenyl, and heteroaryl is 2-, 3-, or 4-
pyridyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 4-
imidazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-
thiazolyl, 1-, 3- or 4-pyrazolyl, 1-, 3- or 4-
~~triazolyl, 2-pyrazinyl, or 1-, 2- or 5-~~
tetrazolyl, each of which is optionally mono-,
di-, or tri-substituted with halogen,
trifluoromethyl, hydroxy, amino, lower
alkylamino, lower dialkylamino, lower
alkylcarboxamido, lower alkyl, lower alkoxy,
with the proviso that 2 adjacent substituents
can together form a 5-7 fused cycloalkyl or
heterocycloalkyl ring;

lower alkanoyl, lower alkylsulfonyl, with the proviso
that R⁴ and R⁵ cannot both be alkanoyl or
alkylsulfonyl; or

NR⁴R⁵ taken together form a C₃-C₆ heterocycloalkyl or
a group of the formula:



wherein e and f are independently 1, 2 or 3 and
the sum of e and f is at least 3; and

G² is


NR⁶ wherein R⁶ is hydrogen or C₁-C₃ alkyl, or

CH(C₀-C₆ alkylene)-G³-R⁷ wherein G³ is CONH,
 CONH(C₁-C₆ alkyl), NH, NH(C₁-C₆ alkyl) and R⁷
 is hydrogen or C₁-C₆ alkyl; or

CONH₂, CO[N(C₁-C₆ alkyl)R⁸] wherein R⁸ is
 hydrogen or C₁-C₆ alkyl;

C₁-C₆ arylalkyl or C₁-C₆ heteroarylalkyl, where
 aryl is phenyl, and heteroaryl is 2-, 3-, or 4-
 pyridyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 4-
 imidazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-
 thiazolyl, 1-, 3- or 4-pyrazolyl, 1-, 3- or 4-
 triazolyl, 2-pyrazinyl, or 1-, 2- or 5-
 tetrazolyl, each of which is optionally mono-,
 di-, or tri-substituted with halogen,
 trifluoromethyl, hydroxy, amino, lower
 alkylamino, lower dialkylamino, lower
 alkylcarboxamido, lower alkyl, lower alkoxy,
 with the proviso that 2 adjacent substituents
 can together form a 5-7 fused cycloalkyl or
 heterocycloalkyl ring;

A² is hydrogen, C₁-C₆ alkyl, (C₁-C₆ alkylene)-G⁴-R⁹
 wherein G⁴ is oxygen or sulfur and R⁹ is hydrogen,
 trifluoromethyl or C₁-C₆ alkyl;




(ii)

wherein heteroaryl is 2-, 3- or 4-pyridyl, 2-, 4- or 5-
 pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-
 oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-
 pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or 1-,

2- or 5-tetrazolyl, each of which is optionally mono- or disubstituted with halogen, trifluoromethyl, amino, C₁-C₆ alkyl, C₁-C₆ alkoxy, with the proviso that tetrazolyl can have at most one substituent;

5 Z^1 is C₁-C₆ alkyl; and

V^2 , Y^2 and A^2 are as defined above;



(iii)

where

Z^2 is carbon or nitrogen;

10 where

when Z^2 is CH, n is 0, 1, 2 or 3 and p is 1, 2, or 3,

R^{10} is carboxamido, or (C₀-C₆ alkylene)-G⁵-R¹¹

wherein G⁵ is NH, NH(C₁-C₆ alkyl) and R¹¹ is hydrogen, C₁-C₆ alkyl, C₁-C₆ arylalkyl or C₁-C₆ heteroarylalkyl, where aryl is phenyl, and heteroaryl is 2-, 3-, or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or 1-, 2- or 5-tetrazolyl, each of which is optionally mono-, di-, or tri-substituted with halogen, trifluoromethyl, hydroxy, amino, lower alkylamino, lower dialkylamino, lower alkylcarboxamido, lower alkyl, lower alkoxy, with the proviso that 2 adjacent substituents can

together form a 5-7 fused cycloalkyl or heterocycloalkyl ring; ;

when Z^2 is carbon, n is 1 or 2 and p is 1 or 2, R^{10} is amino; or

5 when Z^2 is nitrogen, n is 1 or 2 and p is 1 or 2, R^{10} is hydrogen; or

(iv) a nitrogen heterocycle of the formula:



wherein the N-ring represents triazolyl, tetrazolyl, imidazolyl, or pyrazolyl, each of which is optionally substituted with amino, trifluoromethyl, carboxamido, or (C₁-C₆ alkylene)-G⁶-R¹² wherein G⁶ is NH, NH(C₁-C₆ alkyl) and R¹² is hydrogen, C₁-C₆ alkyl, C₁-C₆ arylalkyl or C₁-C₆ heteroarylalkyl, where aryl is phenyl, and heteroaryl is 2-, 3-, or 4-pyridyl, 2-, 4- or 5-pyrimidinyl, 1-, 2- or 4-imidazolyl, 2-, 4-, or 5-oxazolyl, 2-, 4-, or 5-thiazolyl, 1-, 3- or 4-pyrazolyl, 1-, 3- or 4-triazolyl, 2-pyrazinyl, or 1-, 2- or 5-tetrazolyl, each of which is optionally mono-, di-, or tri-substituted with halogen, trifluoromethyl, hydroxy, amino, lower alkylamino, lower dialkylamino, lower alkylcarboxamido, lower alkyl, lower alkoxy, with the proviso that 2 adjacent substituents can together form a 5-7 fused cycloalkyl or heterocycloalkyl ring.

2. A compound according to claim 1, where W is CH, Ar is 2,4,6-trimethylphenyl, R₁ is methyl, and X is N-(2-aminoethyl)-N-(cyclopropylmethyl), N-(2-pyrrolidinoethyl)-N-(cyclopropylmethyl), N-(2-N',N'-dimethylaminoethyl)-N-(propyl), N-(2-(N',N'-dimethyl)aminoethyl)-N-(2-methoxyethyl), N-(2-(N',N'-dimethyl)aminoethyl)-N-(cyclopropylmethyl), N-(2-(N'-ethyl)aminoethyl)-N-(cyclopropylmethyl), N-(2-(N'-methyl-N'-ethyl)aminoethyl)-N-(cyclopropylmethyl), N-(2-(N',N'-diethyl)aminoethyl)-N-(cyclopropylmethyl), N-(2-(N'-methyl)aminoethyl)-N-(cyclopropylmethyl), N-(2-piperidinoethyl)-N-(cyclopropylmethyl), N-(2-pyrrolidinoethyl)-N-(propyl), N-(2-piperidinoethyl)-N-(propyl), N-(2-morpholinoethyl)-N-(cyclopropylmethyl), N-(2-(N'-methyl-N'-ethyl)aminoethyl)-N-(propyl), N-(2-(N'-methyl)piperazinyl)-N-(cyclopropylmethyl), N-(2-(N'-methyl)homopiperazinyl)-N-(cyclopropylmethyl), N-(2-(N'-isopropyl)aminoethyl)-N-(cyclopropylmethyl), N-(3-aminopropyl)-N-(cyclopropylmethyl), N-(2-aminoethyl)-N-(propyl), N-(2-aminoethyl)-N-(ethyl), N-(3-pyrrolidinopropyl)-N-(cyclopropyl methyl), N-(4-pyrrolidinobutyl)-N-(cyclopropyl methyl), N-(2-(N'-2-phenethyl)aminoethyl)-N-(cyclopropylmethyl), N-(2-pyrrolidinoethyl)-N-(2-methoxyethyl), N-(2-pyrrolidinoethyl)-N-(ethyl), N-(2-pyrrolidinoethyl)-N-(butyl), N-(2-pyrrolidinoethyl)-N-(isopropyl), N-(2-(N',N'-dimethyl)aminoethyl)-N-(isobutyl), N-(2-(N'-methyl-N'-ethyl)aminoethyl)-N-(isobutyl), N-(2-(N'-methylpyrrolidin-2-yl)ethyl)-N-(cyclopropyl methyl), N-(2-pyrrolidinoethyl)-N-(isopropyl), N-(2-(N',N'-dimethyl)aminoethyl)-N-(isopropyl), N-(2-(N'-cyclopentyl)aminoethyl)-N-(ethyl), N-(2-(N'-2-(4-methoxy)phenethyl)aminoethyl)-N-(ethyl), N-((1-methyl-2-pyrrolidino)ethyl)-N-(cyclopropyl methyl), N-((1-methyl-2-amino)ethyl)-N-(cyclopropyl methyl).

3. A compound according to claim 1, where W is CH, Ar is 2,4,6-trimethylphenyl, R₁ is methyl, and X is N-(1-oxo-2-(N',N'-dimethyl)aminoethyl)-N-(propyl), N-(2-oxo-2-(N',N'-dimethyl)aminoethyl)-N-(propyl), N-(1-oxo-2-(N',N'-dimethyl)aminoethyl)-N-(cyclopropylmethyl), N-(2-pyrrolidinoethyl)-N-(1-oxo-propyl), N-(2-pyrrolidinoethyl)-N-(1-oxo-cyclopropyl), N-(2-oxo-2-(N',N'-dimethyl)aminoethyl)-N-(cyclopropyl).

4. A compound according to claim 1, where W is CH, Ar is 2,4,6-trimethylphenyl, R₁ is methyl, and X is 2-(S)-(cyclopentylaminomethyl)pyrrolidinyl, 4-(carboxamido)piperidinyl, 3-(cyclopentylamino)pyrrolidinyl, 3-(1-oxo-2-((3-methoxy-4-ethoxy)phenyl)aminoethyl)pyrrolidinyl, 2-(2-(4-methoxy)phenyl)aminoethyl)pyrrolidinyl, 3-(1-oxo-2-((3-methoxy-4-ethoxy)phenyl)ethyl)pyrrolidinyl, 4-(2-((3-methoxy-4-ethoxy)phenyl)aminoethyl)piperidinyl.

5. A compound according to claim 1, where W is N, Ar is 2,4,6-trimethylphenyl, R₁ is methyl, and X is N-(2-pyrrolidinoethyl)-N-(cyclopropylmethyl), N-(2-pyrrolidinoethyl)-N-(propyl), N-(2-pyrrolidinoethyl)-N-(butyl), N-(2-pyrrolidinoethyl)-N-(cyclopropyl), N-(2-(N',N'-dimethyl)aminoethyl)-N-(propyl), N-(2-(N',N'-dimethylaminoethyl)-N-(butyl), N-(2-(N',N'-dimethyl)aminoethyl)-N-(cyclopropylmethyl), N-(2-(N',N'-dimethyl)aminoethyl)-N-(cyclopropyl), N-(2-piperidinoethyl)-N-(propyl), N-(2-piperidinoethyl)-N-(butyl), N-(2-(N',N'-dimethyl)aminoethyl)-N-(cyclobutyl), N-(2-(N',N'-dimethyl)aminoethyl)-N-(cyclopentyl), N-(2-(N',N'-dimethyl)aminoethyl)-N-(2-methoxyethyl), N-(2-(N',N'-

dimethyl)aminoethyl)-N-(isopropyl), N-(2-(N'-ethyl)aminoethyl)-N-(cyclopropyl), piperazinyl

6. A compound according to claim 1, where W is N, Ar is 2,6-dimethyl-4-bromophenyl, R₁ is methyl, and X is N-(2-(N',N'-dimethyl)aminoethyl)-N-(ethyl) or N-(2-pyrrolidinoethyl)-N-(cyclopropyl methyl).


7. A compound according to claim 1, where W is CH, Ar is 2-methyl-4-methoxy phenyl, R₁ is methyl, and X is N-(2-(N',N'-dimethyl)aminoethyl)-N-(cyclopropylmethyl).

8. A compound according to claim 1, where W is CH, Ar is 2,4-dimethylphenyl, R₁ is methyl, and X is N-(2-pyrrolidinoethyl)-N-(cyclopropyl methyl).

9. A compound according to claim 1, where W is CH, Ar is 2,6-dimethylphenyl, R₁ is methyl, and X is N-(2-(N',N'-dimethyl)aminoethyl)-N-(cyclopropylmethyl).

10. A compound according to claim 1, where W is CH, Ar is 2,4,6-trimethylphenyl, R₁ is methyl, and X is N-(2-(1-imidazolyl) aminoethyl)-N-(cyclopropylmethyl), N-(2-(1,2,4-triazol-4-yl)aminoethyl)-N-(cyclopropylmethyl).

11. A compound of the formula:



wherein

each R_a is C_1 - C_6 alkyl;

R_b is hydrogen or methyl;

R_1 is C_1 - C_6 alkyl;

5 R_s is C_1 - C_6 alkyl, $(C_3$ - $C_5)$ cycloalkyl $(C_1$ - $C_3)$ alkyl, $(C_1$ - $C_3)$ alkoxy $(C_1$ - $C_3)$ alkyl, or $(C_3$ - $C_5)$ cycloalkyl;

t is 1, 2 or 3; and

R_x is hydrogen, C_1 - C_6 alkyl, phenyl $(C_1$ - $C_6)$ alkyl where phenyl is optionally mono- or disubstituted independently with C_1 - C_6 alkyl, C_1 - C_6 alkoxy, halogen, or hydroxy; and

10 R_y is hydrogen, C_1 - C_6 alkyl, $(C_3$ - $C_6)$ cycloalkyl; or

NR_xR_y represents pyrrolidinyl, N - $(C_1$ - $C_6)$ alkylpyrrolidin-2-yl, piperidinyl, morpholinyl, or N - $(C_1$ - $C_6)$ alkylpiperazinyl.

12. A compound according to claim 11, wherein R_s is C_1 - C_6 alkyl or cyclopropylmethyl.


13. A compound according to claim 12, wherein R_s is cyclopropyl $(C_1$ - $C_3)$ alkyl.

20 14. A compound according to claim 13, wherein R_x and R_y independently represent hydrogen or C_1 - C_2 alkyl.

15. A compound according to claim 14, wherein each R_a is methyl.

25 16. A compound according to claim 15, where W is nitrogen.

17. A compound of the formula:



wherein

each R_a is C_1 - C_6 alkyl;

R_1 is C_1 - C_6 alkyl;

5 R_s is C_1 - C_6 alkyl, cyclopropyl (C_1 - C_3) alkyl or (C_1 - C_3) alkoxy (C_1 - C_3) alkyl;

t is 1 or 2;

R_x is hydrogen, C_1 - C_6 alkyl, phenyl (C_1 - C_6) alkyl where phenyl is optionally mono- or disubstituted independently with C_1 - C_6 alkyl, C_1 - C_6 alkoxy, halogen, or hydroxy; and

10 R_y is hydrogen, C_1 - C_6 alkyl, (C_3 - C_6) cycloalkyl; or

NR_xR_y represents pyrrolidinyl, N- (C_1 - C_6) alkylpyrrolidin-2-yl, piperidinyl, morpholinyl, or N- (C_1 - C_6) alkylpiperazinyl.

15 18. A compound according to claim 17, wherein R_s is C_1 - C_6 alkyl or cyclopropylmethyl.


19. A compound according to claim 18, wherein R_s is cyclopropyl (C_1 - C_3) alkyl.

20 20. A compound according to claim 19, wherein t is 1 and R_x and R_y independently represent hydrogen or C_1 - C_2 alkyl.

21. A compound according to claim 20, wherein each R_a is methyl.

22. A compound according to claim 21, where W is nitrogen.

23. A compound of the formula:



wherein

each R_a is C_1-C_6 alkyl;

5 R_1 is C_1-C_6 alkyl;

t is 1 or 2;

R_x and R_y are different and represent hydrogen; (C_3 -
 C_7)cycloalkylamino(C_1-C_3)alkyl, carboxamido, (C_3 -
 C_7)cycloalkylamino, C_2-C_6 alkanoyl optionally substituted
 10 in the ω -position with C_1-C_6 alkyl or phenyl optionally
 mono- or disubstituted independently with C_1-C_6 alkyl, C_1 -
 C_6 alkoxy, halogen, or hydroxy, provided that at least one
 of R_x and R_y is hydrogen.

15 24. A compound according to claim 23, wherein R^1 is C_1-C_6
 alkyl and W is nitrogen.

25. A compound according to claim 24, wherein each R_a is
 methyl.

20

26. A compound according to claim 1 which is

4-(N-(2-Pyrrolidinoethyl)-N-cyclopropylmethyl)amino-2-
 methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole;

4-(N-(2-N',N'-Dimethylaminoethyl)-N-propyl)amino-2-
 25 methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole;
 4-(N-(2-N',N'-Dimethylaminoethyl)-N-2-methoxyethyl)amino-
 2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole;

4- (N- (2-N', N'-Dimethylaminoethyl) -N-cyclopropylmethyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

5 4- (N- (2-N' -Ethylaminoethyl) -N-cyclopropylmethyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4- (N- (2- (N' -Ethyl-N' -methyl) aminoethyl) -N-cyclopropylmethyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

10 4- (N-2- (2- (S) -Methoxymethylpyrrolidino) ethyl-N-cyclopropylmethyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4- (N- (2-N', N'-Dimethylaminoethyl) -N-cyclopropylmethyl) amino-2-methyl-9- (4-methoxy-2-methylphenyl) -9H-pyridino[2,3-b]indole;

15 4- (N- (2-N', N'-Diethylaminoethyl) -N-cyclopropylmethyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole; or

4- (N- (2-N' -Methylaminoethyl) -N-cyclopropylmethyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole.

20

27. A compound according to claim 1 which is

4- (N- (2-Piperidinoethyl) -N-cyclopropylmethyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

25 4- (N- (2-Pyrrolidinoethyl) -N-propyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4- (N- (2-Piperidinoethyl) -N-propyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4- (N- (2-Morpholinoethyl) -N-cyclopropylmethyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

30 4- (N- (2-N' -Ethyl-N'-methylaminoethyl) -N-propyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4- (N- (2- (1-Imidazolyl) ethyl) -N-cyclopropylmethyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4 - (N-2 - (N' -Methylpiperazinyl) ethyl -N-cyclopropylmethyl) amino-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4 - (N-2 - (N' -Methylhomopiperazinyl) ethyl -N-cyclopropylmethyl) amino-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4 - (N - (2 -N' -Isopropylaminoethyl) -N-cyclopropylmethyl) amino-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4 - (N - (2 - (2 - (1-Methyl-2-pyrrolidino) ethyl) aminoethyl) -N-cyclopropylmethyl) amino-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole; or

4 - (N - (2 -Piperazinylethyl) -N-cyclopropylmethyl) amino-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole.

28. A compound according to claim 1 which is

4 - (N - (2 -Pyrrolidinoethyl) -N-cyclopropylmethyl) amino-2-methyl-9 - (2,4-dimethylphenyl) -9H-pyridino[2,3-b]indole;

4 - (N - (3 -Pyrrolidinopropyl) -N-cyclopropylmethyl) amino-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4 - (N - (2 -Methyl-2-pyrrolidinoethyl) -N-cyclopropylmethyl) amino-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4 - (N - (4 -Pyrrolidinobutyl) -N-cyclopropylmethyl) amino-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4 - (N - (2 - (4 -Piperidinopiperidinyl) ethyl) -N-cyclopropylmethyl) amino-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4 - (N - (2 - (2 -Phenethylamino) ethyl) -N-cyclopropylmethyl) amino-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4 - (N - (2 -N' -Methylaminoethyl) -N-methyl) amino-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4 - (N - (2 -Pyrrolidinoethyl) -N-2-methoxyethyl) amino-2-methyl-9 - (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4-(N-(2-Pyrrolidinoethyl)-N-ethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole;

4-(N-(2-Pyrrolidinoethyl)-N-butyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole; or

5 4-(N-(2-Pyrrolidinoethyl)-N-2-methylpropyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole.

29. A compound according to claim 1 which is

10 4-(N-(2-N',N'-Dimethylaminoethyl)-N-2-methylpropyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole;

4-(N-(2-(N'-Ethyl-N-methyl)aminoethyl)-N-2-methylpropyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole;

15 4-(N-(2-Pyrrolidinoethyl)-N-isopropyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole;

4-(N-(2-N',N'-Dimethylaminoethyl)-N-isopropyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole.

4-(N-(2-Guanidinoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole;

20 4-(N-(2-(2-(4-Methoxy)phenethylamino)ethyl)-N-ethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole;

4-(N-(2-N'-Cyclopentylaminoethyl)-N-ethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole;

25 4-(N-(2-N',N'-Dimethylaminoethyl)-N-ethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole;

4-(N-(2-N',N'-Dimethylaminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,6-dimethylphenyl)-9H-pyridino[2,3-b]indole; or

30 4-(N-(2-aminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole.

30. A compound according to claim 1 which is

4-(N-(3-Aminopropyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole;

4 - (N - (2 - Amino - 2 - methylethyl) - N - cyclopropylmethyl) amino - 2 - methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H - pyridino[2, 3 - b] indole;

4 - (N - (2 - Aminoethyl) - N - propyl) amino - 2 - methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H - pyridino[2, 3 - b] indole;

5 4 - (N - (2 - Aminoethyl) - N - cyclopropylmethyl) amino - 2 - methyl - 9 - (2, 4 - dimethylphenyl) - 9H - pyridino[2, 3 - b] indole;

4 - (N - (2 - Aminoethyl) - N - cyclopropylmethyl) amino - 2 - methyl - 9 - (2, 6 - dimethylphenyl) - 9H - pyridino[2, 3 - b] indole;

10 4 - (N - (2 - Aminoethyl) - N - ethyl) amino - 2 - methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H - pyridino[2, 3 - b] indole;

4 - (N - (4 - Aminobutyl) - N - cyclopropylmethyl) amino - 2 - methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H - pyridino[2, 3 - b] indole;

4 - (N - (2 - (4 - Triazolyl) ethyl) - N - cyclopropylmethyl) amino - 2 - methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H - pyridino[2, 3 - b] indole;

15 4 - (N - (2 - Carboxamidoethyl) - N - cyclopropylmethyl) amino - 2 - methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H - pyridino[2, 3 - b] indole; or

4 - (N - (2 - N' - acetyl - N' - methylaminoethyl) - N - cyclopropylmethyl) amino - 2 - methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H - pyridino[2, 3 - b] indole.

20

31. A compound according to claim 1 which is

4 - (N - (2 - (N' - Methanesulfonyl - N' - methylamino) ethyl) - N - cyclopropylmethyl) amino - 2 - methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H - pyridino[2, 3 - b] indole;

25 4 - (N - (2 - (N' - Methyl - N' - trifluoromethanesulfonyl) aminoethyl) - N - cyclopropylmethyl) amino - 2 - methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H - pyridino[2, 3 - b] indole;

4 - (N - (2 - (4 - Aminophenylsulfonamido) ethyl) - N - cyclopropylmethyl) amino - 2 - methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H - pyridino[2, 3 - b] indole;

30 4 - (N - (2 - (2 - Thienylsulfonamido) ethyl) - N - cyclopropylmethyl) amino - 2 - methyl - 9 - (2, 4, 6 - trimethylphenyl) - 9H - pyridino[2, 3 - b] indole;

4-(N-(2-(2-(2-Thienyl)-1-oxoethyl)aminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole;

5 4-(N-(2-(2-Phenyl)-1-oxoethyl)aminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole;

4-(N-(2-N',N'-dimethylamino-1-oxoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole;

10 4-(N-(2-N',N'-Dimethylamino-1-oxoethyl)-N-propyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole;

4-(N-(2-N',N'-Dimethylamino-1-oxoethyl)-N-2-methoxyethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole; or

15 4-(N-(2-Pyrrolidino-1-oxoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole.

32. A compound according to claim 1 which is

20 4-(N-(2-N'-Ethylamino-1-oxoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole;

4-(N-(2-(N'-Ethyl-N1-methyl)amino-1-oxoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole;

25 4-(N-(2-(2-(S)-Methoxymethylpyrrolidino)-1-oxoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole;

4-(N-(2-(1-Imidazolyl)-1-oxoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole;

30 4-(N-(2-pyrrolidinoethyl)-N-1-oxopropyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyridino[2,3-b]indole;

4- (N- (2-Pyrrolidinoethyl) -N-cyclopropyloxomethyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4- (N- (2-Dimethylaminoethyl) -N-methoxymethyloxomethyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4- (N- (2-pyridylmethyl) -N-cyclopropylmethyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4- (N- (2-N', N'-Dimethylamino-2-oxoethyl) -N-2-methoxyethyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole; or

4- (N- (2-N', N'-Dimethylamino-2-oxoethyl) -N-propyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole.

4- (N- (2-N', N'-Dimethylamino-2-oxoethyl) -N-2-methoxy-1-oxoethyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4- (N- (2-N', N'-Dimethylamino-1-oxoethyl) -N-cyclopropylmethyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4- (4-Triazolyl) -2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4-Piperazinyl-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4- (4-Methylpiperazinyl) -2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4-Homopiperazinyl-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4- (N- (2- (1-Methyl-2-pyrrolidino) ethyl) -N-cyclopropylcarbonyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4- (N-2- (S) -Pyrrolidinomethylpyrrolidinyl) -2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4- (4-Aminomethylpiperidino) -2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b]indole;

4 - (4-Piperidinopiperidinyl) -2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole;

4 - (4-Carboxamidopiperidinyl) -2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole;

5 4 - (3-Aminopyrrolidinyl) -2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole;

33. A compound according to claim 1 which is

4 - (N- (2- (1-Methyl-2-pyrrolidino) ethyl) -N-cyclopropylmethyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole;

4 - (N-2- (S) -Cyclopentylaminomethylpyrrolidinyl) -2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole;

15 4 - (N-2- (R) -Cyclopentylaminomethylpyrrolidinyl) -2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole;

4 - (N-3-Cyclopentylaminopyrrolidinyl) -2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole;

4 - (N-3- (2- (3-Methoxy-4-ethoxy) phenethyl) amino pyrrolidinyl) -2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole;

4 - (N-3- (1-oxo-2- (3-Methoxy-4-ethoxy) phenethyl) amino pyrrolidinyl) -2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole;

4 - (N-2- (2- (R) - (4-Methoxy) phenethyl) aminomethyl pyrrolidinyl) -2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole;

4 - (N-2- (2- (S) - (4-Methoxy) phenethyl) aminomethyl pyrrolidinyl) -2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole;

30 4 - (N-4- (2- (3-Methoxy-4-ethoxy) phenethyl) aminomethyl piperazinyl) -2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyridino[2,3-b] indole; or

4 - (N- (2-N', N' -Dimethylaminoethyl) -N-cyclopropyl) amino-2-methyl-9- (2,4,6-trimethylphenyl) -9H-pyrimidino[4,5-b] indole.

34. A compound according to claim 1 which is

4-(N-(2-N',N'-Dimethylaminoethyl)-N-methyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole;

4-(N-(2-N',N'-Dimethylaminoethyl)-N-ethyl)amino-2-methyl-9-(4-bromo-2,6-dimethylphenyl)-9H-pyrimidino[4,5-b]indole;

4-(N-(2-N',N'-Dimethylaminoethyl)-N-ethyl)amino-2-methyl-9-(4-chloro-2-methylphenyl)-9H-pyrimidino[4,5-b]indole;

4-(N-(2-Morpholinoethyl)-N-propyl)amino-2-methyl-9-(2-chloro-4-methylphenyl)-9H-pyrimidino[4,5-b]indole;

4-(N-(2-N',N'-Dimethylaminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole;

4-Piperazinyl-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole;

4-(N-(2-Pyrrolidinoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole;

4-(3-Aminopyrrolidinyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole;

4-(2-(S)-Aminomethylpyrrolidino)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole; or

4-(3-Amino-2-triazolyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole.

35. A compound according to claim 1 which is

4-(3-Amino-1-triazolyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole;

4-(3-Amino-4-triazolyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole;

4-(N-(2-Pyrrolidinoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(4-bromo-2,6-dimethylphenyl)-9H-pyrimidino[4,5-b]indole;

4-(N-(2-Pyrrolidinoethyl)-N-propyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole;

4-(N-(2-Pyrrolidinoethyl)-N-butyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole;

4-(N-(2-Pyrrolidinoethyl)-N-cyclopropyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole;

5 4-(N-(2-N',N'-Dimethylaminoethyl)-N-propyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole;

4-(N-(2-N',N'-Dimethylaminoethyl)-N-butyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole;

10 4-(N-(2-Piperidinoethyl)-N-propyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole; or

4-(N-(2-Piperidinoethyl)-N-butyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole.

36. A compound according to claim 1 which is

15 4-(N-(2-N',N'-Dimethylaminoethyl)-N-cyclobutyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole;

4-(N-(2-N',N'-Dimethylaminoethyl)-N-cyclopentyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole;

20 4-(N-(2-N',N'-Dimethylaminoethyl)-N-isopropyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole;

4-(N-(2-N'-Methylaminoethyl)-N-methyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole;

4-(4-(N-Methylamino)piperidino)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole;

25 4-(4-(2-Aminoethyl)piperazinyl)-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole;

4-(N-(2-N'-Ethylaminoethyl)-N-ethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole;

30 (±)-4-(N-(2-N',N'-Dimethylaminoethyl)-N-1-methylpropyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole;

4-(N-(2-N',N'-Diethylaminoethyl)-N-cyclopropylmethyl)amino-2-methyl-9-(2,4,6-trimethylphenyl)-9H-pyrimidino[4,5-b]indole; or

4 - (N - (2 - N' , N' - Diethylaminoethyl) - N - cyclopropyl) amino - 2 - methyl - 9 - (2 , 4 , 6 - trimethylphenyl) - 9H - pyrimidino [4 , 5 - b] indole.

37. A compound according to claim 1 which is

5 4 - (N - (2 - N' , N' - Dimethylaminoethyl) - N - 2 - methoxyethyl) amino - 2 - methyl - 9 - (2 , 4 , 6 - trimethylphenyl) - 9H - pyrimidino [4 , 5 - b] indole;

4 - (N - (2 - N' , N' - Dimethylaminoethyl) - N - 2 - ethoxyethyl) amino - 2 - methyl - 9 - (2 , 4 , 6 - trimethylphenyl) - 9H - pyrimidino [4 , 5 - b] indole;

10 4 - (N - (2 - N' - Ethylaminoethyl) - N - ethyl) amino - 2 - methyl - 9 - (2 - methyl - 4 - chlorophenyl) - 9H - pyrimidino [4 , 5 - b] indole;

4 - (N - (Cyclopentylaminopiperidin - 4 - yl) - N - ethyl) amino - 2 - methyl - 9 - (2 , 4 , 6 - trimethylphenyl) - 9H - pyrimidino [4 , 5 - b] indole;

4 - (N - (2 - N' , N' - Dimethylaminoethyl) - N - isobutyl) amino - 2 - methyl - 9 - (2 , 4 , 6 - trimethylphenyl) - 9H - pyrimidino [4 , 5 - b] indole;

15 4 - (N - (2 - (3 - Methoxy - 4 - ethoxy) phenethyl) aminopiperidin - 4 - yl) - N - ethyl) amino - 2 - methyl - 9 - (2 , 4 , 6 - trimethylphenyl) - 9H - pyrimidino [4 , 5 - b] indole;

20 4 - (N - (S) - (1 - Oxo - 2 - (4 - methoxy) phenethyl) amino pyrrolidin - 2 - yl) - 2 - methyl - 9 - (2 , 4 , 6 - trimethylphenyl) - 9H - pyrimidino [4 , 5 - b] indole; or

4 - (N - (S) - (2 - (4 - Methoxy) phenethyl) amino pyrrolidin - 2 - yl) - 2 - methyl - 9 - (2 , 4 , 6 - trimethylphenyl) - 9H - pyrimidino [4 , 5 - b] indole.

38. A method for the treatment or prevention of
25 physiological disorders associated with an excess of CRF, which method comprises administration to a patient in need thereof a CRF-reducing amount of a compound according to Claim 1.

39. A method of treating affective disorder, anxiety,
30 depression, headache, irritable bowel syndrome, post-traumatic stress disorder, supranuclear palsy, immune suppression, Alzheimer's disease, gastrointestinal diseases, anorexia nervosa or other feeding disorder, drug addiction, drug or alcohol withdrawal symptoms, inflammatory diseases,

cardiovascular or heart-related diseases, fertility problems, human immunodeficiency virus infections, hemorrhagic stress, obesity, infertility, head and spinal cord traumas, epilepsy, stroke, ulcers, amyotrophic lateral sclerosis, hypoglycemia or
5 a disorder the treatment of which can be effected or facilitated by antagonizing CRF, including but not limited to disorders induced or facilitated by CRF, in mammals, comprising: administering to the mammal a therapeutically effective amount of a compound of Claim 1.

10

40. The use of a compound for the manufacture of a medicament for the treatment of affective disorder, anxiety, depression, headache, irritable bowel syndrome, post-traumatic stress disorder, supranuclear palsy, immune suppression,
15 Alzheimer's disease, gastrointestinal diseases, anorexia nervosa or other feeding disorder, drug addiction, drug or alcohol withdrawal symptoms, inflammatory diseases, cardiovascular or heart-related diseases, fertility problems, human immunodeficiency virus infections, hemorrhagic stress,
20 obesity, infertility, head and spinal cord traumas, epilepsy, stroke, ulcers, amyotrophic lateral sclerosis, hypoglycemia or a disorder the treatment of which can be effected or facilitated by antagonizing CRF, including but not limited to disorders induced or facilitated by CRF, in mammals, comprising
25 administering to the mammal a therapeutically effective amount of a compound of Claim 1.

41. A method of treating or preventing a physiological condition in a mammal characterized by the presence of an
30 excess of neuropeptide Y which comprises administering to a mammal in need of such treatment a therapeutically effective amount of a compound according to Claim 1, a prodrug, or a pharmaceutically acceptable salt thereof.

42. A pharmaceutical composition according to Claim 1
for the treatment of disorders or disease states caused by
eating disorders, of obesity, bulimia nervosa, diabetes,
dislipidemia, hypertension, memory loss, epileptic seizures,
5 migraine, sleep disorders, pain, cerebral hemorrhage, shock,
congestive heart failure, nasal congestion or diarrhea.

INTERNATIONAL SEARCH REPORT

I. International Application No

PCT/US 99/07254

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D471/04 C07D487/04 A61K31/435 A61K31/505
 //(C07D487/04, 239:00, 209:00), (C07D471/04, 221:00, 209:00)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,P	WO 98 45295 A (HORVATH RAYMOND F ;NEUROGEN CORP (US); HUTCHINSON ALAN (US)) 15 October 1998 see example II	1-42
Y	WO 95 33750 A (PFIZER ;CHEN YUHPYNG L (US)) 14 December 1995 see abstract; claims 1,13	1-42
Y	WO 96 35689 A (NEUROGEN CORP ;YUAN JUN (US); HUTCHISON ALAN (US)) 14 November 1996 see abstract; claims	1-
Y	WO 94 13676 A (PFIZER ;CHEN YUHPYNG L (US)) 23 June 1994 see abstract; claims 1,11	1-42



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "S" document member of the same patent family

Date of the actual completion of the international search

18 June 1999

Date of mailing of the international search report

15/07/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
 NL - 2280 HV Rijswijk
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl.
 Fax: (+31-70) 340-3016

Authorized officer

Frelon, D

INTERNATIONAL SEARCH REPORT

International Application No
PCT/US 99/07254

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication where appropriate, of the relevant passages	Relevant to claim No.
A	C.E. MÜLLER ET AL.: JOURNAL OF MEDICINAL CHEMISTRY., vol. 39, 1996, pages 2482-2491, XP002106452 WASHINGTON US see page 2488; example II ----	1,40,42
A	K. EGER ET AL.: LIEBIGS ANNALEN DER CHEMIE.,1993, pages 465-470, XP002106453 WEINHEIM DE see page 467; examples 11A,12A ----	1,40,42
A	C.E. MÜLLER ET AL.: JOURNAL OF MEDICINAL CHEMISTRY., vol. 33, 1990, pages 2822-2828, XP002106454 WASHINGTON US see page 2823; table I -----	1,40,42

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 99/07254

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 38,39,41
are directed to methods of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
see further information PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

The compounds of claims 11,17 and 23 as presently figured have not been searched since they do not seem to pertain to the actual invention. This is confirmed by the description on pages 17 to 19 referring to the "preferred compounds of formula I" and by the illustrations of the examples which all are pyridino(2,3-b)indole or pyrimidino(4,5-b)indole derivatives.

INTERNATIONAL SEARCH REPORT

Information on patent family members

I. International Application No

PCT/US 99/07254

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9845295 A	15-10-1998	AU 2622797 A	29-10-1998
WO 9533750 A	14-12-1995	AU 692548 B	11-06-1998
		AU 2453095 A	04-01-1996
		BR 9502708 A	30-04-1996
		CA 2192354 A	14-12-1995
		CN 1150428 A	21-05-1997
		EP 0764166 A	26-03-1997
		FI 964894 A	05-12-1996
		HR 950321 A	28-02-1998
		HU 75774 A	28-05-1997
		JP 9507249 T	22-07-1997
		NO 965237 A	06-02-1997
		NZ 285442 A	27-05-1998
		PL 320631 A	13-10-1997
		ZA 9504677 A	09-12-1996
WO 9635689 A	14-11-1996	US 5804685 A	08-09-1998
		US 5644057 A	01-07-1997
		AU 5679096 A	29-11-1996
		CA 2194756 A	14-11-1996
		EP 0770080 A	02-05-1997
		JP 10506126 T	16-06-1998
		US 5847136 A	08-12-1998
WO 9413676 A	23-06-1994	AT 177101 T	15-03-1999
		AU 690090 B	23-04-1998
		AU 5666494 A	04-07-1994
		CA 2150016 A	23-06-1994
		CN 1097758 A, B	25-01-1995
		CZ 9501584 A	17-01-1996
		DE 69323768 D	08-04-1999
		EP 0674641 A	04-10-1995
		ES 2128544 T	16-05-1999
		FI 935585 A	18-06-1994
		HU 70505 A	30-10-1995
		JP 7509726 T	26-10-1995
		NO 952398 A	16-06-1995
		NZ 258690 A	29-01-1997
		PL 309357 A	02-10-1995
		ZA 9309271 A	12-06-1995